




MALATHION 5 EMULSIVE

An Emulsifiable Liquid An Agricultural Insecticide
Contains 5 lbs. Malathion Per Gallon



GENERAL DIRECTIONS

For use in water as an insecticidal spray. For ground application dilute the recommended amount in 10 to 30 gallons of water per acre; for air application, dilute in 3 to 10 gallons of water per acre, consistent with crop growth and good coverage.

MIXING DIRECTIONS: Pour the required amount of this material through screen into partially-filled spray tank with agitator running while mixing and spraying. If mixture does not mix readily but tends to separate as an oily layer, do not use as injury to plants may result.

USE AND APPLICATION DIRECTIONS

ALFALFA: Aphids (including Yellow Clover Aphid), Potato Leafhoppers, Grasshoppers, and Alfalfa Weevil Larvae - 1½ pints, Armyworms - 2 pints; repeat as needed. Do not apply when alfalfa is in bloom. Do not apply within 7 days of harvest.

BEANS: Mexican Bean Beetle, Leafhopper, Lygus Bugs - 1½ pints; 2 or more applications. Do not apply within 1 day of harvest.

CABBAGE, BROCCOLI, KALE, MUSTARD, TURNIPS: Aphids, Imported Cabbage Worm, Cabbage Looper - 1 to 2 pints; repeat as necessary. Do not apply within 7 days of harvest for cabbage, kale and mustard. Do not apply within 3 days of harvest for broccoli and turnips.

CLOVER: Aphids, Leafhoppers, and Young Grasshoppers - 1½ to 2 pints, repeat as needed. Do not apply when clover is in bloom. Do not apply within 7 days of harvest.

ONIONS: Onion Thrips - 1½ pints; repeat as needed. Make no application on green onions within 3 days of harvest.

PEAS: Pea Aphid - 1½ pints; repeat as needed. Do not apply within 3 days of harvest.

PLUMS, PRUNES: Mealy Plum Louse (Aphid), and Leafhoppers - 1 pint, 1 or more applications. Do not apply within 3 days of harvest.

POTATOES: Aphids - 1 pint, 1 or more applications. Do not apply within 3 days of harvest.

STRAWBERRIES: Aphids - 1½ pints, repeat as needed. Make no application within 3 days of harvest.

TOMATOES: Armyworms, Fruitworms - 2½ quarts, Aphids - 1½ pints, 1 or more applications. (Use 1 pint for aphids; make 1

(continued in column on right)

application.) Do not apply within 5 days of harvest. (In California do not apply within 3 days of harvest.)

ORNAMENTALS: ~~Pine Needles, Oystershell, Juniper, Oak Kermes and Fletcher Scales and Bagworms~~ - dilution per 100 gallons of water: 4 pints for Pine Needle; 1 pint for Oystershell; 2 pints for all others. For scale control, make a thorough full-coverage spray at the time scale crawlers have settled in foliage.

ACTIVE INGREDIENTS	BY WEIGHT
Malathion (0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate)	57%
Aromatic Petroleum Derivative Solvent	35%
INERT INGREDIENTS	8%
	TOTAL 100%

Contains CYTHION the Premium Grade Malathion
EPA Reg. No. 11656-9

CAUTION:

KEEP OUT OF REACH OF CHILDREN

NOTICE OF WARRANTY

WESTERN FARM SERVICE INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

CAUTIONS IN USING

CAUTION! KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS. Harmful if inhaled, swallowed, or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. If material gets in eyes, wash with plenty of water for 15 minutes. Protect from temperatures below 0°F. Avoid contamination of feed and foodstuffs. Wash thoroughly with soap and water after using and before eating and smoking.

NOTE TO PHYSICIANS: Malathion may cause cholinesterase inhibition. Atropine is antidotal.

The Federal Food and Drug Administration has set official tolerances for pesticide residues. Crops having residues in excess of these tolerances should not be marketed without removal of such excess.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

Do not contaminate feed and foodstuffs, drinking fountains, litter and feed troughs. Do not use in milk processing rooms including milk houses and milk storage rooms. Do not use in homes.

Do not re-use empty container. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place. Dispose of empty containers in accordance with local, state or federal recommendations.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment. Residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NET CONTENTS: 5 GALLONS

WESTERN FARM SERVICE, INC., 2401 Crow Canyon Road, SAN RAMON, CA. 94583