

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.



MALATHION 5 EMULSIVE

An Emulsifiable Liquid An Agricultural Insecticide
Contains 5 lbs. Malathion Per Gallon

ACTIVE INGREDIENT:	By Wt.
Malathion (0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate).....	57%
INERT INGREDIENTS.....	43%
TOTAL	100%

Contains CYTHION the Premium Grade Malathion

KEEP OUT OF REACH OF CHILDREN
CAUTION
FOR AGRICULTURAL USE ONLY OR FOR SALE TO USE, AND STORAGE BY SERVICE PERSONS ONLY. DO NOT STORE IN AREAS ACCESSIBLE TO CHILDREN.
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

WESTERN FARM SERVICE, INC.
3075 Citrus Circle, Suite 195
Walnut Creek, California 94598

EPA Reg. No. 11656-9
EPA Est. No. 11656-CA-2

NET CONTENTS: _____ GALLONS

BEST AVAILABLE COPY

GENERAL DIRECTIONS

For use in water as an insecticidal spray. For ground application dilute the recommended amount in 10 to 30 gallons of water per acre; for air application, dilute in 3 to 10 gallons of water per acre, consistent with crop growth and good coverage.

MIXING DIRECTIONS: Pour the required amount of this material through screen into partially-filled spray tank with agitator running while mixing and spraying. If mixture does not mix readily but tends to separate as an oily layer, do not use as injury to plants may result.

USE AND APPLICATION DIRECTIONS

ALFALFA: Aphids (including Yellow Clover Aphid), Potato Leafhoppers, Grasshoppers, and Alfalfa Weevil Larvae - 1 1/2 pints, Armyworms - 2 pints; repeat as needed. Do not apply when alfalfa is in bloom. Do not apply within 7 days of harvest.

BEANS: Mexican Bean Beetle, Leafhopper, Lygus Bugs - 1 1/2 pints; 2 or more applications. Do not apply within 1 day of harvest.

CABBAGE, BROCCOLI, KALE, MUSTARD, TURNIPS: Aphids, Imported Cabbage Worm, Cabbage Looper - 1 to 2 pints; repeat as necessary. Do not apply within 7 days of harvest for cabbage, kale and mustard. Do not apply within 3 days of harvest for broccoli and turnips.

CLOVER: Aphids, Leafhoppers, and Young Grasshoppers - 1 1/2 to 2 pints; repeat as needed. Do not apply when clover is in bloom. Do not apply within 7 days of harvest.

ONIONS: Onion Thrips - 1 1/2 pints; repeat as needed. Make no application on green onions within 3 days of harvest.

PEAS: Pea Aphid - 1 1/2 pints; repeat as needed. Do not apply within 3 days of harvest.

PLUMS, PRUNES: Mealy Plum Louse (Aphid), and Leafhoppers - 1 pint; 1 or more applications. Do not apply within 3 days of harvest.

POTATOES: Aphids - 1 pint; 1 or more applications. Do not apply within 3 days of harvest.

STRAWBERRIES: Aphids - 1 1/2 pints; repeat as needed. Make no application within 3 days of harvest.

TOMATOES: Armyworms, Fruitworms - 2 1/2 quarts. Aphids - 1 1/2 pints; 1 or more applications. (Use 1 pint for aphids; make 1 application.) Do not apply within 5 days of harvest. (In California do not apply within 3 days of harvest.)

ORNAMENTALS: Pine Needle, Oystershell, Juniper, Oak Kermes and Fleascale Scales and Sapporinos - dilution per 100 gallons of water: 4 pints for Pine Needle; 1 pint for Oystershell; 2 pints for all others. For scale control, make thorough full-coverage spray as the time scale crawlers have settled in foliage.

NOTICE OF WARRANTY

WESTERN FARM SERVICE INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION!
Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin, wash thoroughly after handling and before eating or smoking.

NOTE TO PHYSICIAN: Malathion may cause cholinesterase inhibition. Atropine is antidotal.

ENVIRONMENTAL HAZARDS

This product is toxic to shrimp, crab, and fish, keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources.

Highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

REENTRY STATEMENTS

Do not apply this product in such a manner as to directly or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in treated area or in an area about to be treated with this product.

ORAL WARNINGS shall inform workers of areas or fields that may not be entered, without protective equipment or clothing specified on this label, during treatment or until sprays have dried. Follow first aid treatment instructions shown on this label in case of accidental exposure.

REENTRY STATEMENT (Cont'd.)

When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers.

WRITTEN WARNINGS must include the following information.

CAUTION-Area treated with Malathion on _____ **(date)**

Do not enter without appropriate protective clothing until sprays have dried. Follow first aid treatment instructions shown on this product label in case of accidental exposure.

STORAGE & DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

STORAGE

Keep pesticide in original container. Do not use, pour, spill, or store near heat or open flame.

PESTICIDE DISPOSAL

Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. Wastes resulting from the use of this product may be disposed of at an approved waste disposal facility. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Rinse to recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.