

Western Farm Service, Inc.
1969
EPA Reg. No. 148-1100



PARATHION 8

EMULSIFIABLE LIQUID

Contains 8.5% Parathion Per Gallon

ACTIVE INGREDIENT: Parathion 8.5% Per Gallon
INERT INGREDIENTS: 91.5%

KEEP OUT OF REACH OF CHILDREN.

DANGER PELIGRO POISON



NOT FOR HOME USE

FIRST AID TREATMENT
STATEMENTS FOR FIRST AID TREATMENT

EPA Reg. No. 148-1100
EPA Est. No. 148-1100

WESTERN FARM SERVICE, INC.

7775 Center Street
San Diego, CA 92121

ACCEPTED MAY 19 1969

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 148-1100

STATEMENTS FOR FIRST AID TREATMENT

IF SWALLOWED: IF SWALLOWED, DO NOT INDUCE VOMITING. Give 1-2 glasses of water. If unconscious, give water by mouth. Call a doctor immediately.
IF INHALED: If inhaled, get fresh air immediately. If symptoms persist, call a doctor.
IF ON SKIN: Wash with soap and water. Do not use solvents.
IF IN EYES: Flush eyes with water for at least 15 minutes. Call a doctor.

KEEP OUT OF REACH OF CHILDREN.
DANGER - PELOIGRO - POISON
IF SWALLOWED: IF SWALLOWED, DO NOT INDUCE VOMITING. Give 1-2 glasses of water.
IF INHALED: If inhaled, get fresh air immediately.
IF ON SKIN: Wash with soap and water.
IF IN EYES: Flush eyes with water for at least 15 minutes.

REENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.
Reentry into treated fields before expiration of the reentry interval specified on this label is prohibited, unless the protective clothing and equipment specified on this label are used.
Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.
Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product.

ATTENTION

To avoid excessive residues of parathion on food or forage crops always observe the statements found under "Directions for Use" limiting the time before harvest when parathion may be applied.
Because this material is poisonous by skin contact, inhalation or swallowing it should not be used in such a manner or under weather conditions as will permit drift of the spray onto areas not intended to be sprayed. Do not apply or allow drift to areas occupied by unprotected humans or beneficial animals.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
STORAGE AND HANDLING:
1. Handle container carefully so as to avoid damage and prevent leakage.
2. Do not use or store in or around the home.
3. Store container in a well ventilated place.
4. Empty container retains vapor and product residue. Observe all label safeguards until container is cleaned, reconditioned or destroyed.
5. Keep out of reach of children and domestic animals.
DISPOSAL:
Wastes of this pesticide are actually hazardous. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of according to label use instructions, contact your state pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.
Triple rinse empty containers. Flush with water then offer for recycling or reconditioning or puncture to prevent other use and dispose of in an approved landfill or by other procedure approved by state and local authorities.

Vertical text on the right margin: 148-1100

ALFALFA CLOVER, VETCH: Greenbug (aphids), armyworms, grasshoppers - 1/4 to 1/3 pt. per acre. Alfalfa weevil larvae - 1/2 pt. per acre. Do not apply within 15 days of harvest or grazing. Do not reenter treated field within 7 days of application, unless protective clothing and equipment specified on this label are used.

BARLEY, WHEAT, OATS: Greenbug (aphids), armyworms, grasshoppers - 1/4 to 1/3 pt. per acre. Do not apply within 15 days of harvest or grazing. Do not reenter treated field within 3 days of application, unless protective clothing and equipment specified on this label are used.

COTTON: Aphids, red spider mites - 1/3 to 1/2 pt. per acre. Leafhopper - 1/2 pt. per acre. Leaf perforator - 1 pt. per acre. Cabbage looper, stink bugs, salt marsh caterpillar - 1/2 to 1 pt. per acre. Do not hand pick or harvest within 7 days of applications. Do not enter treated field within 3 days of application, unless protective clothing and equipment specified on this label are used.

FRUITS AND NUTS

For control of codling moth, apple maggot, oriental fruit moth, or plum curculio, apply at petal fall and every 10 to 14 days thereafter until control is achieved.

ALMONDS: Aphids - 1/4 pt. per 100 gals. For Oriental Fruit Moth - 1/3 pt. per 100 gals. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. Tent caterpillar, Leafroller - 1/3 to 1/2 pt. per 100 gals. Peach twig borer - 1/2 pt. per 100 gals. Apply for thorough coverage. Do not apply after hulls begin to open. Do not feed treated hulls to livestock. Do not reenter treated field within 6 days of application, unless protective clothing and equipment specified on this label are used.

ALMONDS (Dormant): Parlatoria scale, San Jose scale, Brown Apricot scale, Black scale, overwintering twig borer - 1/2 pt. plus 1-1/2 to 2 gals. Supreme Spray per 100 gals. Do not reenter treated field within 6 days of application, unless protective clothing and equipment specified on this label are used.

APPLES: Codling moth, orange Tortrix moth, San Jose scale, Forbes scale, scurfy scale, woolly apple aphid, mealybug - 1/4 to 1/3 pt. per 100 gals. water or 1 to 1-1/2 pts. per acre. Red banded leafroller - 1/4 to 1/3 pt. per 100 gals. Apply during the prepink or pink stage. Repeat at 7 to 10-day intervals or as needed during the summer months. Eyespotted bud moth, red bug, fruit tree leaf roller, rosy apple aphid, green apple aphid, grasshopper, European red mite, two spotted mite, Pacific mite, Schoene mite, Willamette mite, clover mite - 1/4 pt. per 100 gals. water or 1 pt. per acre. Repeat at 7 to 10 day intervals during summer months. This product may cause injury to fruit and foliage of McIntosh, Jonathan, Cortland, Kendall, McCoun and Melba variety apples. Apply for thorough coverage. Do not apply within 14 days of harvest. Do not reenter treated field within 6 days of application, unless protective clothing and equipment specified on this label are used.

per acre. Aphids, red spider mites - 1/4 pt. per 100 gals. water or 1 to 2 pts. per acre. Apply for thorough coverage. Do not apply within 14 days of harvest. Do not reenter treated field within 6 days of application, unless protective clothing and equipment specified on this label are used.

CITRUS (Oranges, Lemons, Grapefruit): Red scale - 4 to 10 pt. with 1 gal. Supreme Spray or Light Summer Oil in 100 gals. water or 10 pts. per acre. Citricola, cottony cushion scale, yellow scale, purple scale - 1/3 pt. or 1/4 pt. with 1 gal. Supreme Spray or Light Summer Oil in 100 gals. water or 10 pts. per acre. Apply as full coverage spray and restrict use of oil to periods recommended for oil sprays by your State Agricultural Extension Service or Experiment Stations. Citricola, citrus thrips, orange Tortrix moth, Katydid - 1-1/2 pt. per acre. Cottony cushion scale, cutworms - 2 pts. per acre. Fruit tree leaf roller - 1 pt. per acre. Pink scavenger caterpillar - 2-1/2 pts. per acre. Mealybug - 1/3 pt. in 100 gals. water. Workers are not to enter treated areas for any activity requiring substantial contact with treated foliage for 21 days when 4 pints or less per acre per year is applied; 30 days when more than 4 pts. per acre, but no more than 10 pts. per acre per year is applied; 45 days when more than 8 pts. per acre or less than 10 pts. per acre per year is applied. May be applied by aircraft. Do not apply within 30 days of harvest. Western Tussock moth (other than California)

2 pts. per acre. Do not reenter treated field within 21 days of application, unless protective clothing and equipment specified on this label are used.

GRAPES: Mealybug (in California only), leaf folder - 1/4 to 1/3 pt. per 100 gals. water or 1/2 to 3/4 pt. per acre. Red spider mite - 1/4 pt. per 100 gals. water or 1/2 pt. per acre, repeating application at 5 to 7 days. Do not apply within 14 days of harvest. Workers are not to enter treated areas to engage in any activity requiring substantial contact with treated foliage within 21 days following application, unless protective clothing and equipment specified on this label are used.

PEACHES, NECTARINES: Oriental fruit moth, San Jose scale, peach twig borer, cat-facing insects - 1/4 pt. per 100 gals. water or 1 to 1-1/2 pts. per acre. By Air: 1 pt. per acre. Fruit tree leaf roller, cottony peach scale, green peach aphids, red spider mite - 1/4 pt. per 100 gals. water or 1 pt. per acre. (Dormant) San Jose scale, Parlatoria scale, walnut scale, brown apricot scale, mite eggs; overwintering twig borer - 1/2 pt. (not to exceed 2-1/2 pts. per acre) with 1-1/2 to 2 gals. Supreme Spray in 100 gals. water or 1-1/2 pts. with 3 to 4 gals. Supreme Spray per acre. In California, do not apply within 21 days of harvest; for other areas, 14 days. Do not apply more than 5 pts. of PARATHION 8 E.C. per acre between January 1 and harvest. Workers are not to enter treated areas for any activity requiring substantial contact with treated foliage for 21 days after application, unless protective clothing and equipment specified on this label are used.

3063

PARATHION 8 Emulsifiable Liquid

PEARS: Pear leaf blister mite - 1/4 to 1/3 pt. in 100 gals. water in a prebloom spray. By Air: 1 to 1-1/2 pts. per acre. Pear psylla - 1/4 pt. in 100 gals. water as a pre-bloom or post-bloom spray. By Air: 3/4 to 1 pt. per acre. Codling moth, mealy bug, woolly aphid - 1/4 to 1/3 pt. in 100 gals. water or 1-1/2 pts. per acre. By Air: 1-1/2 pts. per acre. Spider mite - 1/4 pt. in 100 gals. water or 1 pt. per acre. By Air: 1 pt. per acre. Under certain conditions injury may result to Bosc pears. Do not apply within 14 days of harvest. Do not reenter treated field within 6 days of application, unless protective clothing and equipment specified on this label are used.

PRUNES, PLUMS: Eye spotted bud moth, mealy plum aphids, Tortrix moth, leaf roller, leaf hopper, aphids, red spider mite - 1/4 to 1/3 pt. per 100 gals. water. By Air: 1 to 1-1/2 pts. per acre. San Jose scale, Parlatoria scale, brown apricot scale, black scale, mealy plum aphid eggs, overwintering twig borer, as a dormant spray - 1/2 pt. in 100 gals. water with 1 gal. Supreme Spray or 1-1/2 pts. with 3 gals. Supreme Spray per acre. Do not apply within 14 days of harvest. Do not reenter treated field within 6 days of application, unless protective clothing and equipment specified on this label are used.

STRAWBERRIES: Leaf roller, red spider mite, lygus bug 1/4 to 1/2 pt. per acre. By Air: 1/2 pt. per acre. Repeat at 7 to 10 day intervals. Strawberry root weevil 1/2 pt. per acre. Do not apply within 14 days of harvest. Do not reenter treated field within 3 days of application, unless protective clothing and equipment specified on this label are used.

VEGETABLES

APTICHOKES: Plum moth, aphid - 1/2 to 1 pt. per acre. Do not apply within 7 days of harvest. Do not reenter treated field within 3 days of application, unless protective clothing and equipment specified on this label are used.

BEANS: Mexican bean beetle, leafhopper, aphids, red spider mites, two spotted spider mite, armyworm, leaf roller, leaf miner - 1/3 to 1/2 pt. per acre. Repeat at 7 to 10 day intervals until controlled. Do not apply within 7 days of harvest. Do not reenter treated field within 3 days of application, unless protective clothing and equipment specified on this label are used.

BEETS: Webworm - 1/2 pt. per acre. Aphids, leafhopper 1/3 to 1/2 pt. per acre. Spider mites - 1/2 pt. per acre. Do not apply within 15 days of harvest or 21 days if tops are to be used for food or feed. Do not reenter treated field within 3 days of application, unless protective clothing and equipment specified on this label are used.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER,

KALE, MUSTARD GREENS: Diamondback moth, imported cabbage worm, armyworms, cabbage loopers - 1/2 pt. per acre. Aphids, thrips - 1/3 to 1/2 pt. per acre. Do not apply within 7 days of harvest to broccoli, Brussels sprouts, or cauliflower or within 10 days to cabbage, kale or mustard greens. Do not reenter treated field within 3 days of application, unless protective clothing and equipment specified on this label are used.

CARROTS: Aphids - 1/2 pt. per acre. Do not apply within 15 days of harvest. Do not use treated carrot tops for food or feed. Do not reenter treated field within 3 days of application, unless protective clothing and equipment specified on this label are used.

CELERY: Celery leafminer, aphids - 1/4 to 1/2 pt. per acre. (1/4 pt. - 15 days to harvest; 1/2 pt. - 21 days; above 1/2 pt. - 30 days.) Do not reenter treated field within 3 days of application, unless protective clothing and equipment specified on this label are used.

CORN: Sap beetles, corn leaf aphid - 1/4 to 1 pt. per acre. Do not apply within 12 days of harvest or cutting for forage. Do not reenter treated field within 6 days of application, unless protective clothing and equipment specified on this label are used.

CUCUMBERS, SQUASH, MELONS: Cucumber beetle, serpentine leaf miner, aphids, spider mite - 1/4 to 1/2 pt. per acre. Leafhopper - 1/2 pt. per acre. Days to harvest: Melons - 7 days; Squash - 15 days; Cucumbers - 15 days. Do not harvest within the indicated days of application. For cucumbers do not apply before vining. Do not reenter treated field within 3 days of application, unless protective clothing and equipment specified on this label are used.

LETTUCE: Aphids, cabbage loopers, armyworms, leaf miner - 1/3 to 1/2 pt. per acre. Do not apply to head lettuce within 7 days nor to leaf and bibb lettuce within 21 days of harvest. Do not reenter treated field within 3 days of application, unless protective clothing and equipment specified on this label are used.

ONIONS: Thrips - 1/3 pt. per acre. Repeat at weekly intervals as necessary. Onion maggots - 1/2 pt. per acre in furrow at time of planting. Do not apply within 15 days of harvest. Do not reenter treated field within 3 days of application, unless protective clothing and equipment specified on this label are used.

PEAS: Aphids, pea weevil - 1/4 pt. per acre. By Air: 1/3 pt. per acre. Do not apply within 10 days of harvest. Do not reenter treated field within 3 days of application, unless protective clothing and equipment specified on this label are used.

PEPPERS: Aphids, serpentine leaf miner - 1/3 pt. per acre. By Air: - 3/4 pt. per acre. Do not apply within 15 days of harvest. Do not reenter treated field within 3 days of application, unless protective clothing and equipment specified on this label are used.

POTATOES: Flea beetle, leafhoppers, serpentine leafminer, grasshopper, whitefly, aphid - 1/4 pt. per acre. By Air: - 1/3 pt. per acre. Potato psyllid - 1/3 to 1 pt. per acre. Do not apply within 5 days of harvest. Do not reenter treated field within 3 days of application, unless protective clothing and equipment specified on this label are used.

SPINACH: Aphids - 1/4 to 1/2 pt. per acre. By Air: 1/2 pt. per acre. Do not apply within 14 days of harvest. Do not reenter treated field within 3 days of application, unless protective clothing and equipment specified on this label are used.

TOMATOES: Flea beetle, leafhopper, serpentine leaf

miner, grasshopper, white flies, aphids 1/2 pt. per acre by air. Do not apply within 10 days of harvest. Do not reenter treated field within 3 days of application, unless protective clothing and equipment specified on this label are used.

TURNIPS: Diamond-back moth, imported cabbage worm, armyworms - 1/4 to 1/3 pt. per acre. By Air: 1/3 pt. per acre. Aphids, thrips - 1/4 pt. per acre. By Air: 1/3 pt. per acre. Do not apply within 10 days of harvest. Do not reenter treated field within 3 days of application, unless protective clothing and equipment are used.

EPA Reg. No. 11656-14-AA

EPA Est. No. 11656-CA-2

WESTERN FARM SERVICE, INC.
3075 Citrus Circle, Suite 195
Walnut Creek, CA 94598