



# PARATHION 4 AN EMULSIFIABLE LIQUID

Contains 4 lbs. Parathion per Gal.

(PARATHION MIXTURE, LIQUID —  
POISONOUS LIQUID, CLASS B)

ACTIVE INGREDIENTS	By Weight
Parathion (0,0-diethyl 0-p-nitrophenyl phosphorothioate)	47.6%
Aromatic Petroleum Derivative Solvents	46.7%
INERT INGREDIENTS	5.7%
TOTAL	100.0%

EPA Reg. No. 11656-15 AA NET CONTENTS 30 GALS.

**STOP — READ LABEL BEFORE USING!**

NOT FOR HOME USE

(Red) **POISON** (Red) **DANGER**  
 See Side Panels for Antidote & Precautions  Keep Out of Reach of Children  
 CAN KILL YOU

### FISH AND WILDLIFE PRECAUTIONS

**POISONOUS TO FISH AND WILDLIFE.** This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal or wastes.

Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your State Agricultural Extension Service or Experiment Station.

**Use Only When Wearing the Following Protective Equipment and Clothing:**

1. Wear water-proof pants, coat, hat, rubber boots or rubber overshoes.
2. Wear safety goggles.
3. Wear a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provisions of 30CFR Part 11.
4. Wear heavy-duty, natural rubber gloves.

### WORK SAFETY RULES

Keep all unprotected persons and children away from treated area or where there is danger of drift. **DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.**

Do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to Parathion and describe his condition. After first aid is given (see First Aid Treatment Section) and if a doctor cannot come, take victim to clinic or hospital.

**IMPORTANT!** Before removing gloves, wash them with soap and water. Always wash hands, face and arms with soap and water before smoking, eating or drinking.

**AFTER WORK,** take off all clothing. Shower, using soap and water. Wash and change leaving job. Do not wear contaminated clothing and protective equipment after each use. Respirator should be cleaned according to instructions included.

**TO PREVENT PERSONAL INJURIES:** Keep all persons and animals out of treated areas. Treated areas should not be used for insecticide and volatile residues in any manner other than recommended indoors, provide mechanical exhaust.

Do not apply or allow to drift in treated humans or beneficial animals. Do not apply parathion on food or forage crops unless found under "Directions for Use" in harvest when parathion may be applied.

Do not use or store in or around reach of children and domestic animals or feed products. Bury spillage; clean solution.

### POISON SIGNS

Parathion is a very dangerous body on contact with all skin surfaces with this material must be removed immediately. Some of the signs and symptoms are: headache, nausea, vomiting, cramps, watery eyes, tightness in chest, labored breathing, sweating, watering of eyes, drooling, nose, muscle spasms and coma.

### FIRST AID TREATMENT

Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to Parathion and describe his condition.

**IF BREATHING HAS STOPPED** immediately and maintain until doctor arrives.

**IF SWALLOWED** and victim is conscious, induce vomiting by giving vomit quickly. Induce vomiting by giving soapy or strong salty water.

### DANGER — POISON — PRECAUTIONS

#### CAN KILL YOU IF SWALLOWED

This product can kill you if swallowed even in small amounts: spray mist or dust may be fatal if swallowed.


#### CAN KILL YOU BY SKIN CONTACT

This product can kill you if touched by hands or spilled or splashed on skin, in eyes or on clothing (liquid goes through clothes).

#### CAN KILL YOU IF BREATHED

This product can kill you if vapors, spray mist or dust are breathed.

Red



WFS 15-01 8-75

MANUFACTURED IN U.S.A. BY **WESTERN FARM SERVICE, INC.** 2401 Crow Canyon Road, San

RD  
ACE

# ION 4 LIQUID

ion per Gal.  
E, LIQUID —  
CLASS B)

	By Weight
.....	47.6%
vents .....	46.7%
.....	5.7%
TOTAL	100.0%

T CONTENTS 30 GALS.

BEFORE USING!

USE

(Red)  
**DANGER**

Keep Out of  
Reach of  
Children

### FISH AND WILDLIFE PRECAUTIONS

**POISONOUS TO FISH AND WILDLIFE.** This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal or wastes.

Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your State Agricultural Extension Service or Experiment Station.

#### Use Only When Wearing the Following Protective Equipment and Clothing:

1. Wear water-proof pants, coat, hat, rubber boots or rubber overshoes.
2. Wear safety goggles.
3. Wear a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provisions of 30CFR Part 11.
4. Wear heavy-duty, natural rubber gloves.

### WORK SAFETY RULES

Keep all unprotected persons and children away from treated area or where there is danger of drift. **DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.**

Do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to Parathion and describe his condition. After first aid is given (see First Aid Treatment Section) and if a doctor cannot come, take victim to clinic or hospital.

**IMPORTANT!** Before removing gloves, wash them with soap and water. Always wash hands, face and arms with soap and water before smoking, eating or drinking.

**AFTER WORK,** take off all work clothes and shoes. Shower, using soap and water. Wear only clean clothes when leaving job. Do not wear contaminated clothing. Wash protective clothing and protective equipment with soap and water after each use. Respirator should be cleaned and filter replaced according to instructions included with respirator.

**TO PREVENT PERSONAL INJURY AND POSSIBLE FATALITIES:** Keep all persons and animals out of treated areas for 48 hours. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not use in any manner other than recommended on this label. If handled indoors, provide mechanical exhaust ventilation.

Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. To avoid excessive residues of parathion on food or forage crops, always observe the statements found under "Directions for Use," limiting the time before harvest when parathion may be applied.

Do not use or store in or around the home. Keep out of reach of children and domestic animals. Do not store near food or feed products. Bury spillage; clean up area with strong lye solution.

### POISON SIGNS (Symptoms)

Parathion is a very dangerous poison. It rapidly enters the body on contact with all skin surfaces and eyes. Clothing wet with this material must be removed immediately. Exposed persons must receive prompt medical treatment or they may die.

Some of the signs and symptoms of poisoning are: Headache, nausea, vomiting, cramps, weakness, blurred vision, pinpoint pupils, tightness in chest, labored breathing, nervousness, sweating, watering of eyes, drooling or frothing of mouth and nose, muscle spasms and coma.

### FIRST AID TREATMENT

Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to Parathion and describe his condition.

**IF BREATHING HAS STOPPED,** start artificial respiration immediately and maintain until doctor sees victim.

**IF SWALLOWED** and victim is awake (conscious) make him vomit quickly. Induce vomiting by sticking finger down throat or by giving soapy or strong salty water to drink. Repeat until vomit

is clear. Never give anything by mouth to an unconscious person. Have victim lie down and keep quiet. See doctor immediately.

**IN CASE OF CONTACT,** immediately flush eyes or skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. See doctor immediately.

### NOTE TO PHYSICIAN

**Antidote —** Administer atropine sulfate in large doses, 2 to 4 mg. intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 10-minute intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. **DO NOT GIVE MORPHINE OR TRANQUILIZERS.** Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement; **VERY CLOSE SUPERVISION IS INDICATED FOR AT LEAST 48 HOURS.**

**POST TREATED AREA.** Consult your State Agricultural Extension Service or Experiment Station regarding posting treated areas.

**DISPOSAL OF EMPTY CONTAINER —** Do not re-use this container. Completely empty the contents and bury the unused chemical at least 18 inches deep in an isolated location away from water supplies. Rinse out the inside of the container with water to which has been added detergent and caustic soda. Carefully discard the rinse solution by burying at least 18 inches deep in an isolated area away from water supplies. Puncture and crush empty metal container and bury at least 18 inches deep in a supervised public or private dump.

**CROP USE DIRECTIONS —** Use only according to attached label instructions.

### NOTICE OF WARRANTY

**WESTERN FARM SERVICE, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED,** concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

**WESTERN FARM SERVICE, INC. 2401 Crow Canyon Road, San Ramon, CA. 94583**

WESTERN



FARM SERVICE

# PARATHION 4

## AN EMULSIFIABLE LIQUID

Contains 4 lbs. Parathion per Gal.

ACTIVE INGREDIENTS	By Weight
Parathion (0,0-diethyl 0-p-nitrophenyl phosphorothioate) .....	47.6%
Aromatic Petroleum Derivative Solvents .....	46.7%
INERT INGREDIENTS .....	5.7%
TOTAL	100.0%

EPA Reg. No. 11656-15 AA                      NET CONTENT                      GALS.

**STOP — READ LABEL BEFORE USING!**

NOT FOR HOME USE

**POISON**

See Side Panels  
for Antidote &  
Precautions



CAN KILL YOU

**DANGER**

Keep Out of  
Reach of  
Children



Red

### DANGER—POISON—PRECAUTIONS

#### CAN KILL YOU IF SWALLOWED

This product can kill you if swallowed even in small amounts; spray mist or dust may be fatal if swallowed.



#### CAN KILL YOU BY SKIN CONTACT

This product can kill you if touched by hands or spilled or splashed on skin, in eyes or on clothing (liquid goes through clothes).



#### CAN KILL YOU IF BREATHED

This product can kill you if vapors, spray mist or dust are breathed.

### NOTICE OF WARRANTY

WESTERN FARM SERVICE, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

**WESTERN FARM SERVICE, INC.**

2401 Crow Canyon Road  
San Ramon, California 94583

WFS 15-01S 8-75

AC

## GENERAL DIRECTIONS

Read entire label. Use strictly in accordance with cautions, warnings and directions, and with applicable state and federal regulations.

Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers.

Keep cool in shade — Cut of direct sunlight.

Wash empty container with alkali and destroy by burying in a safe place away from water supplies and run-off. Never re-use.

This product is intended to be applied only by specified and trained operators of aircraft or power-driven ground equipment. This product is not to be used around homes or home gardens. USE INDICATED AMOUNT PER 100 GALLONS WATER FOR CONVENTIONAL GROUND SPRAYERS UNLESS OTHERWISE SPECIFIED.

GROUND APPLICATION: Vegetable and Field Crops — 30 to 250 gals.; Cotton — 3 to 20 gals.; Deciduous Fruit Crops — 60 to 800 gals. (On Young Fruit Plantings — Use a minimum of 15 gals. water per acre).

AIR APPLICATION: Field Crops — 3 to 10 gals.; Orchard Crops — 10 to 20 gals.

## USE AND APPLICATION DIRECTIONS

### FIELD AND FORAGE CROPS

**COTTON:** Aphids, Red Spider Mites, Leafhoppers —  $\frac{3}{4}$  pt. per acre in water to cover. For Red Spider Mites, apply when mites appear or when leaves begin turning yellow to reddish brown. Repeat once or twice at 5 to 7 day intervals. Leaf Perforator — 1 qt. per acre in water to cover. Cabbage Looper, Stink Bugs, Salt Marsh Caterpillars — 1 pt. per acre in water to cover. Do not hand pick or harvest within 7 days of application.

**RICE:** Tadpole Shrimp, Leafminer — Apply  $\frac{1}{5}$  pt. per acre by air in water to cover. Hold water in check for 3 days after application. Do not apply within 1 day of harvest.

**TOBACCO (Field):** Aphids, Suckly —  $\frac{1}{2}$  pt. per acre. Make no application within 5 days of priming or 15 days of cutting. Avoid contact with plant juices when priming or cutting tobacco. When necessary to enter treated fields within 24 hours after application, protective clothing should be worn. By Air:  $\frac{3}{4}$  pt. per acre in water to cover. (Seedbed): Aphids —  $\frac{1}{2}$  pt. per acre.

**VETCH, ALFALFA, CLOVER, WHEAT, OATS, BARLEY:** Greenbug (Aphids), Armyworms, Grasshoppers —  $\frac{1}{2}$  to  $\frac{2}{3}$  pt. per acre in water to cover. By Air:  $\frac{1}{2}$  to  $\frac{2}{3}$  pt. per acre in water to cover. Alfalfa Weevil Larvae —  $\frac{2}{3}$  pt. per acre. Apply when alfalfa is 12 to 14 inches long. For severe infestation, repeat application to stubble after first cutting. Do not apply within 15 days of harvest. Do not graze animals on treated pasture or cut for hay within 15 days of application. Birds and other wildlife in treated areas may be killed.

### VEGETABLES

**ARTICHOKE:** Plume Moth — 2 pts. per acre in water to cover by ground or air. Aphids — 1 pt. per acre. Do not apply within 7 days of harvest.

**BEANS:** Mexican Bean Beetle, Leafhopper, Aphids, Red Spider Mites, Two-Spotted Spider Mite, Armyworms, Leaf Roller, Leaf Miner —  $\frac{1}{2}$  pt. per acre. By Air: 1 pt. per acre in water to cover. Repeat at 7 to 10-day intervals until controlled. Do not apply within 15 days of harvest.

**BEET:** Webworm —  $\frac{2}{3}$  pt. per acre. By Air: 1 pt. per acre in water to cover. Aphids, Leafhoppers —  $\frac{1}{2}$  pt. per acre. By Air:  $\frac{3}{4}$  to 1 pt. per acre in water to cover. Spider Mites — 1 pt. per acre. Do not apply within 15 days of harvest or 21 days if tops are to be used for food or feed.

**CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KALE, MUSTARD GREENS:** Diamondback Moth, Imported Cabbage Worm, Armyworm —  $\frac{1}{2}$  to 1 pt. per acre. By Air:  $\frac{3}{4}$  to 1 pt. per acre in water to cover. Specific Aphid and Thrip Control —  $\frac{1}{2}$  pt. per acre. By Air:  $\frac{3}{4}$  to 1 pt. per acre in water to cover. Do not apply to broccoli, brussels sprouts and cauliflower within 7 days of harvest and to cabbage, kale and mustard greens within 10 days of harvest.

**CARROTS:** Aphids —  $\frac{1}{2}$  pt. per acre, applying 300 gals. diluted spray per acre. By Air:  $\frac{3}{4}$  pt. per acre in water to cover. Do not use treated carrot tops for food or feed. Do not apply within 15 days of harvest.

**CELERY:** Celery Leafminer, Aphids —  $\frac{1}{2}$  to 1 pt. per acre. By Air:  $\frac{3}{4}$  to 1 pt. per acre in water to cover. Do not apply within 21 days of harvest.

**CORN:** Corn Rootworm (Adults) —  $\frac{1}{2}$  pt. per acre in water to cover. Make full-coverage applications to foliage when adult beetles become abundant. Repeat as necessary. Do not apply within 12 days of harvest or cutting for forage.

**CUCUMBER, SQUASH, MELONS:** Cucumber Beetle, Melonworm, Pickleworm, Serpentine Leaf Miner, Aphids —  $\frac{1}{2}$  pt. per acre. By Air: Cucumbers, Melons —  $\frac{3}{4}$  pt. per acre in water to cover. For Melonworm and Pickleworm, apply at first bloom or when insects appear. Squash —  $\frac{2}{3}$  pt. per acre in water to cover. Do not apply to cucumbers before vining or within 15 days of harvest. Do not apply to melons within 7 days of harvest and to squash within 15 days of harvest.

**LETTUCE:** Aphids, Loopers, Armyworms, Leaf Miner —  $\frac{3}{4}$  to 1 pt. per acre in water to cover. Do not apply to head lettuce within 21 days of harvest.

**ONIONS:** Thrips —  $\frac{2}{5}$  to 1 pt. per acre. Repeat at weekly intervals as necessary. By Air:  $\frac{3}{4}$  pt. per acre in water to cover. Do not apply within 15 days of harvest. Onion Maggots — 1 pt. per acre in water to cover. Apply in the furrow at the time of planting onion seeds.

**PEAS:** Aphid, Pea Weevil —  $\frac{1}{2}$  pt. per acre in water to cover. By Air:  $\frac{3}{4}$  pt. per acre in water to cover. Do not apply within 10 days of harvest.

**PEPPERS:** Aphids —  $\frac{1}{2}$  to 1 pt. per acre. Serpentine Leaf Miner —  $\frac{1}{3}$  pt. per acre. By Air:  $\frac{3}{4}$  pt. per acre in water to cover. Do not apply within 15 days of harvest.

**POTATOES:** Colorado Potato Beetle, Flea Beetle, Leafhopper, Serpentine Leaf Miner, Grasshoppers, Whitefly, Aphids, Potato Psyllid —  $\frac{2}{3}$  pt. per 100 gals. water (200 diluted gals. per acre). By Air:  $1\frac{1}{2}$  pts. per acre in water to cover. Do not apply within 5 days of harvest.

**SPINACH:** Aphids —  $\frac{1}{2}$  to 1 pt. per acre in water to cover. By Air: 1 pt. per acre in water to cover. Do not apply within 14 days of harvest.

**TURNIPS:** Diamondback Moth, Imported Cabbage Worm, Armyworms —  $\frac{1}{2}$  to  $\frac{2}{3}$  pt. per acre. By Air:  $\frac{3}{4}$  pt. per acre in water to cover. Specific Aphid and Thrip Control —  $\frac{1}{2}$  pt. per acre. By Air:  $\frac{3}{4}$  pt. per acre in water to cover. Do not apply within 10 days of harvest.

**PRE-PLANTING SOIL TREATMENT:** Lettuce, Cabbage, Cauliflower, Carrots, Beans — Symphylids (Centipedes), Springtails — 5 qts. per acre in water to cover. Work into top 3 to 6 inches of soil. Keep all persons and animals out of treated area for 48 hours. Do not contaminate ponds or streams.

## FRUITS AND NUTS

**ALMONDS (Dormant):** Parlatoria (Olive) Scale, San Jose Scale, Brown Apricot Scale, Black Scale, Mite Eggs, Overwintering Twig Borer — 1 pt. plus  $1\frac{1}{2}$  to 2 gals. VOLCK Supreme Spray or 3 to 4 gals. KLEENUP Flowable Emulsion.

**APPLES:** Codling Moth, Plum Curculio, Grange Tortrix, San Jose, Forbes and Scruffy Scales, Red-Banded Leaf Roller (Second and Third Brood), Woolly Apple Aphid, Mealybug —  $\frac{1}{2}$  to  $\frac{2}{3}$  pt. By Air: 3 pts. per acre in water to cover. Bud Moth, Red Bug, Fruit Tree Leaf Roller, Rosy Apple Aphid, Green Apple Aphid, Red-Banded Leaf Roller (First Brood), Grasshoppers, European Red Mite, Two-Spotted Mite, Pacific Mite, Schoenus Mite, Willamette Mite, Clover Mite —  $\frac{1}{2}$  pt. By Air: 2 pts. per acre in water to cover. Repeat at 7 to 10-day intervals during summer months. This product may cause injury to fruit and foliage of McIntosh Apples and related varieties. Do not apply within 14 days of harvest.

**APRICOTS:** Codling Moth, Oriental Fruit Moth, Tortrix —  $\frac{2}{3}$  pt. Bud Moth, Fruit Tree Leaf Roller, Red Spider Mites, Aphids —  $\frac{1}{2}$  pt. San Jose Scale, Brown Apricot Scale, Black Scale, Mite Eggs, Overwintering Twig Borer (In Dormant) — 1 pt. (not over 5 pts. per acre) plus  $1\frac{1}{2}$  to 2 gals. VOLCK Supreme Spray or 3 to 4 gals. KLEENUP Flowable Emulsion. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. Do not apply within 14 days of harvest.

**CHERRY:** Plum Curculio, Cherry Fruitworm, Oriental Fruit Moth, Bud Moth, Tortrix, Cankerworm —  $\frac{1}{2}$  pt. By Air: 2 pts. per acre in water to cover. Aphids, Red Spider Mites:  $\frac{1}{2}$  to  $\frac{1}{2}$  pt. By Air: 1 to  $1\frac{1}{2}$  pts. per acre in water to cover. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. Do not apply within 14 days of harvest.

**GRAPES:** Leaf Folder —  $\frac{1}{2}$  to  $\frac{2}{3}$  pt. Mealybug — 1 to 3 pts. per acre. By Air: 1 to  $1\frac{1}{2}$  pts. per acre in water to cover. Red Spider Mites:  $\frac{1}{2}$  to 2 pt. per acre. Repeat at 5 to 7 days. By Air:  $\frac{3}{4}$  pt. per acre in water to cover. Do not apply within 14 days of harvest. Do not enter vineyards within 21 days following treatment.

**PEACHES, NECTARINES:** Plum Curculio, Oriental Fruit Moth, Cat-Facing Insects —  $\frac{1}{2}$  pt. By Airplane: 2 pts. per acre in water to cover. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. Apply when crawlers appear. Fruit Tree Leaf Roller, Cottony Peach Scale, Green Peach Aphid, Red Spider Mites —  $\frac{1}{2}$  pt. By Airplane:  $1\frac{1}{2}$  pts. per acre in water to cover. Peach Twig Borer — 1 pt. San Jose Scale, Olive Scale, Walnut Scale, Brown Apricot Scale, Overwintering Twig Borer (In Dormant) — 1 pt. (not over 5 pts. per acre) plus  $1\frac{1}{2}$  to 2 gals. VOLCK Supreme Spray or 3 to 4 gals. KLEENUP Flowable Emulsion. California Area — Do not apply within 21 days of harvest. Do not apply more than once after bloom. Do not apply more than 5 lbs. actual parathion per acre between Jan. 1 and harvest. All Other Areas — Do not apply within 14 days of harvest. Do not apply more than 5 lbs. actual parathion per acre per year. Do not enter orchards within 21 days following treatment.

**PEARS:** Pearleaf Blister Mite —  $\frac{1}{2}$  to  $\frac{2}{3}$  pt. in pre-blossom spray. 2 to 3 pts. per acre in water to cover. Pear Psylla —  $\frac{1}{2}$  pt. either in pre-blossom or post-blossom spray. By Air:  $1\frac{1}{2}$  to 2 pts. per acre in water to cover. Use dosages listed under APPLE for control of Codling Moth, Mealybug, Woolly Aphid and certain Spider Mites. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. Do not apply within 14 days of harvest. Under certain conditions injury may result to Bosc Pears.

**PRUNE, PLUM:** Plum Curculio —  $\frac{1}{4}$  to  $\frac{2}{3}$  pt. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. Bud Moth, Mealy Plum Louse, Tortrix, Leaf Roller, Leafhoppers, Aphids, Red Spider Mites —  $\frac{1}{2}$  to  $\frac{2}{3}$  pt. By Air: 2 to 3 pts. per acre in water to cover. San Jose Scale, Parlatoria Scale, Brown Apricot Scale, Black Scale, Mealy Plum Aphid Eggs, Mite Eggs, Overwintering Twig Borer (In Dormant) — 1 pt. (not over 5 pts. per acre) plus  $1\frac{1}{2}$  gals. VOLCK Supreme Spray or 3 to 4 gals. KLEENUP Flowable Emulsion. Do not apply within 14 days of harvest.

**STRAWBERRY:** Leaf Roller, Red Spider Mite, Lygus Bugs, Whiteflies —  $\frac{1}{2}$  to  $\frac{2}{3}$  pt. per acre. Repeat at 7 to 10-day intervals. By Air:  $\frac{3}{4}$  pt. per acre in water to cover. Strawberry Root Weevil — 1 pt. Do not apply within 14 days of harvest.

**TOMATOES:** Flea Beetle, Leafhopper, Serpentine Leaf Miner, Grasshoppers, Whitefly, Aphids —  $\frac{1}{2}$  to 1 pt. per acre with regular fungicide program, using 200 gals. diluted spray per acre. By Air: 1 to 2 pts. per acre in water to cover. Do not apply within 10 days of harvest.

## ORNAMENTALS

**Commercial Plantings Only — Not to be Used Adjacent to Residences or Residential Areas:** Aphids, Whiteflies, Red Spider Mites, Leafhoppers —  $\frac{1}{2}$  pt. per acre. Repeat against spider mites and leafhoppers in 7 to 10 days. Mealybugs, Scales, Leaf Tiers —  $\frac{2}{3}$  pt. per acre against young or crawler stages. Repeat application may be necessary in 3 weeks.