

COASTOX CARBARYL CUTWORM BAIT

(Contains 5% Sevin®)

COASTOX CARBARYL CUTWORM BAIT is a special pelletized formulation for the control of cutworms, armyworms, and crickets.

GENERAL USE DIRECTIONS

This product may be applied by hand equipment, ground equipment or aircraft. Apply when air is calm to avoid drift to adjacent crops.

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

RECOMMENDATIONS

ALFALFA, SORGHUM: Cutworm, Crickets. Use 30 lbs. per acre. Do not apply within 3 days of harvest on alfalfa. Do not use within 21 days of harvest on sorghum or grains.

ASPARAGUS: Cutworm. Use 40 lbs. per acre. Apply during cutting season. Do not repeat application within three days. Do not apply within 1 day of harvest.

BEETS: Cutworm, Armyworm. Use 40 lbs. per acre. Do not apply within 3 days of harvest. Do not apply within 14 days if tops are to be used for food or feed.

BEANS: Cutworm, Armyworm. Use 40 lbs. per acre. No time limitations.

BROCCOLI, BRUSSEL SPROUTS, CABBAGE, CAULIFLOWER, HEAD LETTUCE: Cutworm, Armyworm. Use 40 lbs. per acre. Do not apply within 3 days of harvest.

CARROTS: Cutworm. Use 40 lbs. per acre. No time limitations.

CORN: Cutworm. Use 40 lbs. per acre. No time limitations.

CHINESE CABBAGE, COLLARDS, ENDIVE (ESCAROLE), KALE, LEAF LETTUCE, MUSTARD GREENS, PARSLEY, SPINACH, SWISS CHARD, SUGAR BEETS: Cutworm, Armyworm. Use 40 lbs. per acre. Do not use within 14 days of harvest.

COTTON: Cutworm. Use 30 lbs. per acre. Do not apply within 7 days of harvest.

EGGPLANT, PEPPERS, TOMATOES: Cutworms. Use 40 lbs. per acre. No time limitations.

GRAPES: Cutworm. Use 40 to 48 lbs. per acre. No time limitations.

PARSNIPS, RADISHES, RUTABAGA, KOHLRABI: Cutworm. Use 40 lbs. per acre. Do not use within 3 days of harvest.

CUCUMBER, MELONS, SQUASH: Cutworm. Use 20 lbs. per acre. No time limitations.

STRAWBERRIES: Cutworm. Use 40 lbs. per acre. Do not apply within 1 day of harvest.

ORNAMENTAL PLANTS, SHRUBS, AND FLOWERS: Armyworms, Crickets, Cutworm, Sowbug. Use 40 lbs. per acre. For Millipedes use 175 lbs. per acre. Bait should be applied to the ground around the shrubs and trees. Do not apply to ivy as injury may result.

PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals

CAUTION!

May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. Change contaminated clothing daily. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinestase inhibitor. Atropine is antidote.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

Avoid drift of this product to any crop for which this chemical is not registered. Avoid drift to any crop for which use may be registered but closer to harvest than the recommended time limits for this chemical. Avoid drift to any feed or forage crop.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS.

REENTRY STATEMENT

Do not enter treated areas without protective clothing until dusts have settled.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product.

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-methylcarbamate) 5.0%

INERT INGREDIENTS..... 95.0%

100.0%

• Copyright Union Carbide Corporation, U. S. Pat. No. 2,903,478

KEEP OUT OF REACH OF CHILDREN

CAUTION

FOR AGRICULTURAL USE ONLY OR FOR SALE TO USE, AND STORAGE BY SERVICE PERSONS ONLY. DO NOT STORE IN AREAS ACCESSIBLE TO CHILDREN.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE

Keep in original container. Store in cool, dry place. Protect from excessive heat.

For help with any spill, leak, fire or exposure involving this material, call day or night 800-424-9300.

DISPOSAL

Wastes resulting from the use of this product may be disposed of at an approved waste disposal facility.

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTE OF WARRANTY

WESTERN FARM SERVICE MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USES EXTENDING BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORD WITH THE STATEMENTS MADE ON THIS LABEL.

EPA Reg. No. 11656-21

EPA Est. No. 11656-CA-03

NET CONTENTS _____ POUNDS

WESTERN FARM SERVICE, INC.
3075 Citrus Circle, Suite 515
Walnut Creek, CA 94598

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended for the pesticide registered under EPA Reg. No. 11656-21

JUN 28 1985

ACCEPTED