

SEP 21 1987

Ms. Gail Shinn  
Western Farm Service, Inc.  
3075 Citrus Circle, Suite 195  
Walnut Creek, CA 94598

Dear Ms. Shinn:

Subject: Additional Brand Name  
Coastox Carbaryl Cutworm Bait  
EPA Registration No. 11656-21  
Your Application Dated June 23, 1987

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

At your next printing of labeling for this product add a Statement of Practical Treatment to your label to read as listed below:

**STATEMENT OF PRACTICAL TREATMENT**

**IF IN EYES:** Flush with plenty of water. Get medical attention if irritation persists.

**IF ON SKIN OR CLOTHING:** Remove contaminated clothing. Wash skin with soap and warm water. Get medical attention if irritation persists.

**IF INHALED:** Remove to fresh air if effects occur and apply artificial respiration if indicated.

**IF SWALLOWED:** Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Call a physician.

15873:I/C:Edwards:E-15:KENCO:9/15/87:9/25/87:CB:vo:ek:rw:

OFFICE FILE COPY

-2-

We have adjusted our records to include the additional brand name listed below.

Additional Brand Name

Carbaryl Cutworm Bait

It is understood that the label will comply with the most recently accepted primary label for which this registration was issued.

Sincerely yours,

Dennis H. Edwards, Jr.  
Product Manager (12)  
Insecticide-Rodenticide Branch  
Registration Division (TS-767C)

Enclosure



## CARBARYL CUTWORM BAIT

(Contains 5% Sevin<sup>®</sup>)

### ACTIVE INGREDIENT:

Carbaryl (1-Naphthyl N-methylcarbamate) .....	5.0%
INERT INGREDIENTS .....	95.0%
	100.0%

\* Copyright Union Carbide Corporation,  
U. S. Pat. No. 2,903,478

### PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals

#### CAUTION

May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. Change contaminated clothing daily. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

**NOTE TO PHYSICIAN:** Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

#### ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

Avoid drift of this product to any crop for which this chemical is not registered. Avoid drift to any crop for which use may be registered but closer to harvest than the recommended time limits for this chemical. Avoid drift to any feed or forage crop.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

### REENTRY STATEMENT

Do not enter treated areas without protective clothing until dusts have settled.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product.

KEEP OUT OF REACH OF CHILDREN

### CAUTION!

FOR AGRICULTURAL USE ONLY OR FOR SALE TO, USE, AND STORAGE BY SERVICE PERSONS ONLY. DO NOT STORE IN AREAS ACCESSIBLE TO CHILDREN.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

#### STORAGE

Keep in original container. Store in cool, dry place. Protect from excessive heat.

For help with any spill, leak, fire or exposure involving this material, call day or night 800-424-9300.

#### DISPOSAL

Wastes resulting from the use of this product may be disposed of at an approved waste disposal facility.

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### NOTICE OF WARRANTY

WESTERN FARM SERVICE MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

EPA Reg. No. 11656-21  
EPA Est. No. 11656-CA-001

NET CONTENTS

WESTERN FARM SERVICE, INC.

3075 Citrus Circle, Suite 195  
Walnut Creek, CA 94598

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

11656-21

COASTOX CARBARYL pelletized formulation myworms, and crickets.

**GENERAL USE DIRECTIONS:** This product may be applied to ground equipment or to avoid drift to adjacent crops.

Subsequent planting of soil must be made within 14 days of label(s).

**RECOMMENDATIONS:** ALFALFA, SORGHUM, etc. per acre. Do not apply to alfalfa. Do not use within 14 days of harvest.

ASPARAGUS: Cutworm during cutting season. Do not apply within three days. Do not apply to tops.

BEETS: Cutworm. Do not apply within 3 days of harvest. Do not apply to tops.

BEANS: Cutworm. No time limitations.

BROCCOLI, BRUSSELS SPOURGE, CAULIFLOWER, etc. Do not apply within 14 days of harvest.

CARROTS: Cutworm. No time limitations.

CORN: Cutworm. No time limitations.

CHINESE CABBAGE (ESCAROLE), KALE, GREENS, PARSLEY, SUGAR BEETS: Cutworm. Do not use within 14 days of harvest.

COTTON: Cutworm. Do not apply within 7 days of harvest.

EGGPLANT, PEPPER, etc. 40 lbs. per acre. No time limitations.

GRAPES: Cutworm. No time limitations.

PARSNIPS, RADISH, etc. Cutworm. Use 40 lbs. per acre. No time limitations.

CUCUMBER, MELON, etc. 40 lbs. per acre. No time limitations.

STRAWBERRIES: Do not apply within 14 days of harvest.

ORNAMENTAL PLANTS: Armyworms, Crickets, etc. per acre. For Millipedes should be applied to trees. Do not apply to lawns.



## CARBARYL CUTWORM BAIT

(Contains 5% Sevin®)

### ACTIVE INGREDIENT:

Carbaryl (1-Naphthyl N-methylcarbamate) .....	5.0%
INERT INGREDIENTS .....	95.0%
	100.0%

\* Copyright Union Carbide Corporation,  
U. S. Pat. No. 2,903,478

KEEP OUT OF REACH OF CHILDREN

### CAUTION!

FOR AGRICULTURAL USE ONLY OR FOR  
SALE TO, USE, AND STORAGE BY SERVICE  
PERSONS ONLY. DO NOT STORE IN AREAS  
ACCESSIBLE TO CHILDREN.

SEE SIDE PANEL FOR ADDITIONAL  
PRECAUTIONARY STATEMENTS.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage  
or disposal.

#### STORAGE

Keep in original container. Store in cool, dry place. Pro-  
tect from excessive heat.

For help with any spill, leak, fire or exposure involving  
this material, call day or night 800-424-9300.

#### DISPOSAL

Wastes resulting from the use of this product may be  
disposed of at an approved waste disposal facility.

Completely empty bag into application equipment. Then  
dispose of empty bag in a sanitary landfill or by incinera-  
tion, or, if allowed by State and local authorities, by  
burning. If burned, stay out of smoke.

### NOTICE OF WARRANTY

WESTERN FARM SERVICE MAKES NO WARRANTY OF  
MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR  
OTHERWISE, EXPRESS OR IMPLIED, concerning this pro-  
duct or its uses which extend beyond the use of the product  
under normal conditions in accord with the statements made  
on this label.

EPA Reg. No. 11656-21

EPA Est. No. 11656-CA-01

NET CONTENTS

WESTERN FARM SERVICE, INC.

3075 Citrus Circle, Suite 195

Walnut Creek, CA 94598

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

11656-21

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans & Domestic Animals

#### CAUTION

May be harmful if swallowed. Avoid breathing  
dust. Avoid contact with eyes, skin or clothing.  
Wash thoroughly after handling. Change con-  
taminated clothing daily. Do not apply this pro-  
duct in such a manner as to directly or through  
drift expose workers or other persons. The area  
being treated must be vacated by unprotected  
persons.

**NOTE TO PHYSICIAN:** Carbaryl is a moderate,  
reversible cholinesterase inhibitor. Atropine is  
antidotal.

#### ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic in-  
vertebrates. Do not apply directly to water or  
wetlands. Do not contaminate water by cleaning  
of equipment or disposal of wastes.

Avoid drift of this product to any crop for which  
this chemical is not registered. Avoid drift to any  
crop for which use may be registered but closer  
to harvest than the recommended time limits for  
this chemical. Avoid drift to any feed or forage  
crop.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in  
a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORD-  
ANCE WITH PRECAUTIONARY STATEMENTS AND  
DIRECTIONS.

### REENTRY STATEMENT

Do not enter treated areas without protective clothing  
until dusts have settled.

Because certain states may require more restrictive  
reentry intervals for various crops treated with this pro-  
duct, consult your State Department of Agriculture for  
further information.

Written or oral warnings must be given to workers who  
are expected to be in a treated area or in an area about  
to be treated with this product.

COASTOX CARBARYL CUTWORM BAIT is a special  
pelletized formulation for the control of cutworms, ar-  
myworms, and crickets.

### GENERAL USE DIRECTIONS

This product may be applied by hand equipment,  
ground equipment or aircraft. Apply when air is calm  
to avoid drift to adjacent crops.

Subsequent planting of food and feed crops in treated  
soil must be made with crops on this or other carbaryl  
label(s).

### RECOMMENDATIONS

ALFALFA, SORGHUM: Cutworm, Crickets. Use 30 lbs.  
per acre. Do not apply within 3 days of harvest on  
alfalfa. Do not use within 21 days of harvest on sorghum  
or grains.

ASPARAGUS: Cutworm. Use 40 lbs. per acre. Apply  
during cutting season. Do not repeat application within  
three days. Do not apply within 1 day of harvest.

BEETS: Cutworm, Armyworm. Use 40 lbs. per acre. Do  
not apply within 3 days of harvest. Do not apply within  
14 days if tops are to be used for food or feed.

BEANS: Cutworm, Armyworm. Use 40 lbs. per acre.  
No time limitations.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE,  
CAULIFLOWER, HEAD LETTUCE: Cutworm, Ar-  
myworm. Use 40 lbs. per acre. Do not apply within 3  
days of harvest.

CARROTS: Cutworm. Use 40 lbs. per acre. No time  
limitations.

CORN: Cutworm. Use 40 lbs. per acre. No time  
limitations.

CHINESE CABBAGE, COLLARDS, ENDIVE  
(ESCAROLE), KALE, LEAF LETTUCE, MUSTARD  
GREENS, PARSLEY, SPINACH, SWISS CHARD,  
SUGAR BEETS: Cutworm, Armyworm. Use 40 lbs. per  
acre. Do not use within 14 days of harvest.

COTTON: Cutworm. Use 30 lbs. per acre. Do not ap-  
ply within 7 days of harvest.

EGGPLANT, PEPPERS, TOMATOES: Cutworms. Use  
40 lbs. per acre. No time limitations.

GRAPES: Cutworm. Use 40 to 48 lbs. per acre. No time  
limitations.

PARSNIPS, RADISHES, RUTABAGA, KOHLRABI:  
Cutworm. Use 40 lbs. per acre. Do not use within 3 days  
of harvest.

CUCUMBER, MELONS, SQUASH: Cutworm. Use 20  
lbs. per acre. No time limitations.

STRAWBERRIES: Cutworm. Use 40 lbs. per acre. Do  
not apply within 1 day of harvest.

ORNAMENTAL PLANTS, SHRUBS, AND FLOWERS:  
Armyworms, Crickets, Cutworm, Sowbug. Use 40 lbs.  
per acre. For Millipedes use 175 lbs. per acre. Bait  
should be applied to the ground around the shrubs and  
trees. Do not apply to ivy as injury may result.

BEST AVAILABLE