

11656-21

12/6/2001

**FIRST CHOICE**

1/3

ACCEPTED  
DEC 6 2001**CARBARYL CUTWORM BAIT**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticides  
registered under  
EPA Reg. No. 11656-21

(Contains 5% Sevin®) N-Methyl Carbamate

For the control of cutworms, armyworms and crickets

**READ ENTIRE CONTAINER LABEL BEFORE USING THIS PRODUCT****ACTIVE INGREDIENT:**

Carbaryl (1-Naphthyl N-Methylcarbamate)	5.0%
<b>INERT INGREDIENTS</b>	95.0%
<b>TOTAL</b>	100.0%

EPA REG. NO. 11656-21

EPA EST. NO. 11656-CA-02

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

For Agricultural use only or for sale to, use, and storage by service persons only. Do not store in areas accessible to children. See side panel for additional precautionary statements.

**FIRST AID****N-Methyl Carbamate**

**If Swallowed:** Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If In Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses. If present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If On Skin Or Clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If Inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**NOTE TO PHYSICIAN:** Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS & DOMESTIC ANIMALS****CAUTION**

May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing.

**PERSONAL PROTECTIVE EQUIPMENT**

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions of cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is extremely toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning equipment or disposal of wastes. Avoid drift of this product to any crop for which this chemical is not registered. Avoid drift to any crop for which use may be registered but closer to harvest than the recommended time limits for this chemical. Avoid drift to any feed or forage crop.

**GENERAL INFORMATION**

FIRST CHOICE CARBARYL CUTWORM BAIT is a special pelletized formulation for the control of cutworms, armyworms, crickets, darkling ground beetles, grasshoppers, mole crickets and sowbugs. This product may be applied by hand equipment, ground equipment or aircraft. Apply when air is calm to avoid drift to adjacent crops. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, Waterproof gloves, Shoes plus socks.

**FIELD AND VEGETABLE CROPS**

CROP	LBS BAIT/ ACRE	PHI	SPECIFIC DIRECTIONS
Asparagus	40	1	Max. seasonal application of 120 lbs./A. to spears and 80 lbs./A to ferns. Retreatment interval 3 days when applied to spears and 7 days when applied to ferns after harvest. Total number of applications is 3 prior to harvest with 2 additional applications to ferns only. Total maximum seasonal application for ferns and spears is 200 lbs.
Beans, fresh and dried (Snap, navy & kidney) Pea, fresh & dried: cowpea, lentil	40	3 Succulents 21 Dried 14 Forage 21 Hay	Maximum seasonal application rate of 120 lbs./A/crop
Garden beets (roots), carrots, horseradish, radish, parsnip, rutabaga, salsify, turnips (roots)	40	7	Maximum seasonal application rate of 120 lbs./A/crop. Retreatment interval = 7 days. Total number of applications/crop can be 6 (some of which must be at lower than maximum single application rate).
Sugar Beets	30	28	PHI for roots and forage. Maximum seasonal application rate of 60 lbs./A/crop. Retreatment interval = 14 days
Mustard Greens	40	14	Maximum seasonal application rate of 120 lbs./A/crop. Retreatment interval = 7 days. Total number of applications/crop can be 4 (some of which must be at a lower than maximum single application rate).
Brussels sprouts, broccoli, cabbage, cauliflower, Chinese cabbage, collards, kale, kohlrabi	40	3 - Brussels sprouts, broccoli, cabbage, kohlrabi, cauliflower  14 - Chinese cabbage, kale, collards	Maximum seasonal application rate of 120 lbs./A/season. Retreatment interval = 7 days. Total number of applications/crop can be 4 (some of which must be at lower than maximum single application rate).
Corn (field and pop)	40	14 - Forage, silage 48 - Grain, fodder	Maximum seasonal application rate of 160 lbs./A/crop. Retreatment interval = 14 days
Corn (sweet)	40	2 - Ears 14 - Forage 48 - Fodder	Maximum seasonal application rate of 320 lbs./crop. Retreatment interval = 3 days
Cucumber, melon, squash	20	3	Maximum seasonal application rate of 120 lbs./A/crop. Retreatment interval = 7 days
Lettuce (head & leaf), endive, dandelion, parsley, spinach, Swiss chard, Hanover salad, rhubarb, garden beets (tops), turnips (tops)	40	14 - 15	Maximum seasonal application rate of 120 lbs./A/crop. Retreatment interval = 7 days. Total number of applications/crop can be 5 (some of which must be at lower than maximum single application rate).
Grain sorghum (not sweet)	40	14 - Grazing Forage 21 - Grain, fodder	Maximum seasonal application rate of 120 lbs./A/crop. Retreatment interval = 7 days. Total number of applications /crop can be 4 (some of which must be at lower than maximum single application rate).
Strawberry	40	7	Maximum seasonal application rate of 200 lbs./A/crop. Retreatment interval = 7 days
Tomato, pepper, eggplant	40	3	Maximum seasonal application rate of 160 lbs./A/crop. Retreatment interval = 7 days. Total number of applications/crop can be 7 (some of which must be at lower than maximum single application rate).
Ornamental plants, shrubs, and flowers	40	N/A	Armyworms, crickets, cutworm, and sowbug. Bait should be applied to the ground around the shrubs and trees. Do not apply to ivy as injury may result.
	175		For control of millipedes, use the higher rate.

**STORAGE AND DISPOSAL**

Do not contaminate, water, food or feed by storage or disposal.

**STORAGE**

Keep in original container. Store in cool, dry place. Protect from excessive heat. For help with any spill, leak, fire or exposure involving this material, call day or night, (800) 424-9300.

**DISPOSAL**

Wastes resulting from the use of this product may be disposed of at an approved waste disposal facility. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**NOTICE OF WARRANTY**

WESTERN FARM SERVICE MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED, concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

NET WEIGHT: 50 lbs.(22.68 kgs.)

Manufactured by:  
**WESTERN FARM SERVICE, INC.**  
P. O. Box 1168  
Fresno CA 93715

FIRST CHOICE® is a registered trademark of Western Farm Service, Inc.

SEVIN® is a registered trademark of Rhône-Poulenc. U.S. Pat. No. 2,903,478