

CENTER PANEL

EL BIANCO BRAND

PERTHANE 4-E

Emulsifiable Liquid
(Contains 4 lb. Perthane per gal.)

ACTIVE INGREDIENTS:

DIETHYL DIPHENYL DICHLOROETHANE-----	45.8%
RELATED REACTION PRODUCTS-----	2.4%
INERT INGREDIENTS-----	51.8%
	<u>100.0%</u>

CAUTION

KEEP OUT OF REACH OF CHILDREN. →
AVOID BREATHING SPRAY MIST. →
AVOID CONTACT WITH SKIN. IF SKIN CONTACTED, WASH OFF WITH SOAP &
WATER.
WASH WITH SOAP AND WATER AFTER HANDLING.
AVOID CONTAMINATION OF FEED AND FOODSTUFFS.

Do Not Use Or Store Near Heat Or Open Flame.

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS AND ANTIDOTE.

NET CONTENTS _____ GALLON

U.S.D.A. Reg. No. 4110

WHITE CHEMICAL COMPANY
1310 W. WATKINS ROAD PHOENIX, ARIZONA

LEFT

OBSERVE THE FOLLOWING USE PRECAUTIONS

WARNING

Harmful if swallowed. Avoid skin contact with solution. In case of skin contact, wash with soap and water. Avoid breathing spray mist. Avoid storage near feed or food products. Do not store near fire or flame.

ANTIDOTE: In case of skin contact, wash thoroughly with soap and water. For eyes, flush with water and obtain medical attention promptly.

INTERNAL: Give a teaspoonful of salt in a glass of warm water, and repeat until vomit fluid is clear. Keep victim prone and quiet. Call a physician at once.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified.

Consult state agricultural extension service for additional information, as the timing, number, and rate of applications needed will vary with local conditions.

CAUTION: Do not apply where material will drift on grazing dairy cattle, or animals being finished for slaughter. This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply the product only as specified on label.) TT

DIRECTIONS FOR USE

Place recommended amount of this material into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing, and spraying operation. Do not allow mixture to stand. If mixture does not mix readily but tends to separate as an oily layer, do not use as injury to plants may result. Do not combine emulsifiable liquids with wettable powders in the same spray tank unless previous use of the materials being combined has proved physically compatible.

WARNING
KEEP OUT OF REACH OF CHILDREN
AND DOMESTIC ANIMALS.

ACCEPTED

2-24-69

411-52

RECOMMENDATIONS

For ground application, dilute in 10 to 30 gallons of water, and for air application, dilute in 3 to 10 gallons of water, consistent with crop growth.

COLE CROPS (Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kohlrabi) Cabbage looper, imported cabbage worm. Apply 1 quart Perthane 4-E per acre in sufficient water to provide good coverage. Apply as soon as insects appear. Repeat applications at 7-10 day intervals or as needed to achieve control. Do not apply later than 3 days of harvest.

LETTUCE Cabbage looper, imported cabbage worm, leafhoppers -- Apply 1 quart Perthane 4-E per acre in sufficient water to provide good coverage. Apply as soon as insects appear. Repeat applications at intervals of 7-10 days, or as needed to maintain control. Do not apply later than 4 days of harvest.

SPINACH Cabbage looper, leafhoppers -- Apply 1 quart Perthane 4-E per acre in sufficient water to provide good coverage. Apply as soon as insects appear. Repeat at intervals of 7-10 days or as needed to maintain control. Do not apply later than 7 days of harvest.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

2-24-69

4110-52