



CAPTAN SULFUR 10-50 DUST

(FUNGICIDE-MITICIDE)

ACTIVE INGREDIENTS	By Weight
*Captan	10%
Sulfur	49%
INERT INGREDIENTS	41%
TOTAL	100%

*N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide.

EPA REG. NO. 11656-43 AA NET WEIGHT LBS.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS. AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

NOTE: Sulfur may cause severe fruit and foliage injury to certain crops. Consult your State Agricultural Experiment Station or your State Agricultural Extension Service for additional information consistent with these label cautions and directions. Sulfur may burn foliage when temperature is high. Do not make sulfur applications at such times. Do not use within 2 weeks of an oil spray treatment. When growing crops for processing, consult the processor before applying sulfur.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

STORE IN COOL, DRY PLACE. PROTECT FROM EXCESSIVE HEAT.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

CAUTION: KEEP OUT OF REACH OF CHILDREN

CAUTION: Avoid inhalation of dust. Avoid contact with skin. Do not store or transport near feed or food. Do not apply under conditions involving possible drift to food, forage, or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes.

WESTERN FARM SERVICE, INC.

2401 Crow Canyon Road
San Ramon, California 94583

WFS 43-01 8-75

DIRECTIONS FOR USE

This product is formulated for application by ground or air equipment.

CELERY: Pink Rot, Damp-Off: (Field) — 20 lbs. per acre. (Seed Bed) — 35 to 40 lbs. per acre. Apply when all seeds have germinated. Repeat as necessary.

CUCUMBERS: Botrytis Rot (Bunch Rot), Powdery Mildew, Silver Mite, Storage Rots — 45 to 50 lbs. per acre. Make 2 applications before bloom and 1 application just after bloom. Repeat periodically making up to 3 cover applications before bunches close.

PEACHES: Brown Rot, Storage Rots, Powdery Mildew, Silver Mite, Peach Scab — 50 lbs. per acre. Make 2 to 3 prebloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make preharvest application where necessary.

PLUMS: Brown Rot, Powdery Mildew, Silver Mite — 45 lbs per acre. Make 2 to 3 prebloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make preharvest application where necessary.

POTATOES: Early and Late Blight — Begin applications when plants are 2 to 4 inches high and repeat at 5 to 7-day intervals as long as disease threatens. Apply 25 to 30 lbs. per acre.

ROSES (Field): Rust, Powdery Mildew — 35 to 40 lbs. per acre. Apply to give even coverage of upper and lower leaf surfaces at first sign of disease. Repeat at regular intervals as necessary to maintain control.

SOIL TREATMENT: Pre-planting for Damp-Off on Vegetable Field Crops — 35 lbs. per acre. Dust or broadcast, then disc into first few inches of soil surface.

SPINACH: Downy Mildew, Powdery Mildew — 35 to 40 lbs. per acre. Apply when all seeds have germinated. Repeat at 7 to 10-day intervals as long as diseases are active.

SQUASH (California Only): Powdery Mildew, Downy Mildew, Angular Leaf Spot, Anthracnose — Apply 30 lbs. per acre on young plants increasing to 35 lbs. per acre on mature plants. Apply at first sign of disease. Repeat at 5 to 10-day intervals until danger of infection is past. Do not apply to squash varieties susceptible to sulfur injury.

STRAWBERRIES: Botrytis Rot, Powdery Mildew — 30 lbs. per acre immediately after picking. Repeat applications at 7-day intervals through picking and after harvest.

TOMATOES (Established Plants): Early Blight, Late Blight, Anthracnose, Russet Mite — On young plants, apply 25 to 30 lbs. per acre, increasing to 35 to 40 lbs. per acre on mature plants. Apply at first sign of insects and diseases. Repeat at 5 to 7-day intervals as necessary for control. Because of the sulfur content of this dust, first obtain permission from your canner or processor before using it on tomatoes intended for sale to such canner or processor.

NOTICE OF WARRANTY

WESTERN FARM SERVICE MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its use which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

ACCEPTED WITH COMMENTS

WARNING: May be fatal if swallowed! Do not get in eyes or on skin. Do not breathe dust or spray mist. Coarse sprays are less likely to drift. Wash thoroughly after handling. Keep children and domestic animals off treated areas. Do not graze treated areas or feed livestock treated forage. Keep away from children and domestic animals.

ANTIDOTE

EXTERNAL: If spilled on skin, wash off immediately with plenty of soap and cold water. If in eyes, flush out with water and call doctor immediately.

INTERNAL: If swallowed, give raw white of egg. Induce vomiting with a mixture of salt, soap or mustard in warm water. Call doctor at once.

NON WARRANTY: Vineland Chemical Company and seller warrant that this material will conform to ingredients indicated on label, but make no other warranty implied or expressed. Neither Vineland Chemical Company nor seller shall be held responsible for personal injury, property damage or other loss resulting from the handling, storage or use of the product, whether or not in accordance with directions. Buyer assumes all risks and liability therefrom and accepts and uses the product on these conditions. No person is authorized to waive this notice.

NET CONTENT

GALLONS

*A
Turf
Fungicide*



VI-CAD

A Liquid Cadmium Turf Fungicide

ACTIVE INGREDIENT:

Cadmium Chloride 20.1%

INERT INGREDIENT:

..... 79.9%

Total 100.0%

Total Cadmium (Cd) 12.3%

DANGER **POISON** **DANGER**

CAUTION: Keep Out of Reach of Children
See Antidote statement and additional cautions
on left side panel.

Manufactured By

VINELAND CHEMICAL COMPANY

VINELAND, NEW JERSEY

USDA Reg. No. 2853-10

VC-518
Litho in U.S.A.

ACCEPT
June 19,
UNDER THE FEDERAL IN-
FUNGICIDE AND RODENT
FOR ECONOMIC POISON
ED U.S. DEPT. OF AGRICULTURE
TO AGRICULTURAL COMMENT