

Handwritten notes and stamps in the top left corner, including a date stamp and some illegible text.



MAY CAUSE SKIN IRRITATION      HARMFUL IF SWALLOWED

Avoid contact with skin or clothing. Avoid inhalation. In case of contact, flush skin with plenty of water; for eyes, flush with plenty of water for 15 minutes. Get medical attention. Wash contaminated clothing before reuse.

This product is non-selective and is harmful to all vegetation. Spray, drift, seepage and drainage will injure or kill all vegetation.

**NOTICE OF WARRANTY**

WESTERN FARM SERVICE, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE OF OTHERWISE, EXPRESS OR IMPLIED concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

**COTTON DEFOLIANT 1**

Contains 2 Pounds Sodium Chlorate per Gallon

<b>ACTIVE INGREDIENT</b>	<b>By Weight</b>
Sodium Chlorate (NaClO <sub>3</sub> ).....	19.7%
<b>INERT INGREDIENTS</b> .....	<u>80.3%</u>
<b>TOTAL</b>	<b>100.0%</b>

EPA Reg. No. 11656-44 AA

**NET CONTENTS:      GALS.**

**CAUTION:**  
**KEEP OUT OF REACH OF CHILDREN**

WFS 44-01 8-75

MANUFACTURED IN U.S.A. BY **WESTERN FARM SERVICE, INC. 2401 Crow Canyon Road, San Ramon, CA**

Best defoliation of content of the plant are deroliant late to insure percent and preferably time of application. You least 30 days old. The r ment Station or State A

FOR AIRPLANE A DEFOLIANT in 10-15 ga TION: Apply 2-2½ gal water per acre. The def leaf to obtain best resul Satisfactory defoliation with dense cotton a seco

Do not apply within feed gin waste to livest

Wash spray equipr must be thoroughly wa original container. Do n

Do not contaminate wastes. Apply this prod

Do not reuse empty by perforating or crush supplies.



# COTTON DEFOLIANT 1

Contains 2 Pounds Sodium Chlorate per Gallon

<b>ACTIVE INGREDIENT</b>	<b>By Weight</b>
Sodium Chlorate (NaClO <sub>3</sub> ) .....	19.7%
<b>INERT INGREDIENTS</b> .....	<u>80.3%</u>
<b>TOTAL</b>	<b>100.0%</b>

EPA Reg. No. 11656-44 AA

**NET CONTENTS: GALS.**

**CAUTION:**  
**KEEP OUT OF REACH OF CHILDREN**

WFS 44-01 8-75

## DIRECTIONS FOR USE

Best defoliation occurs on cotton when the nitrogen and moisture content of the plant are nearly or completely exhausted. It is better to apply defoliant late to insure a higher yield of better quality fiber. At least 75 percent and preferably 85 to 90 percent of the bolls should be open at the time of application. Youngest bolls expected to make cotton should be at least 30 days old. The recommendations of your State Agricultural Experiment Station or State Agriculture Extension Service should be followed.

**FOR AIRPLANE APPLICATION:** Apply 2-2½ gallons of COTTON DEFOLIANT in 10-15 gallons of water per acre. **FOR GROUND APPLICATION:** Apply 2-2½ gallons of COTTON DEFOLIANT in 15-30 gallons of water per acre. The defoliant must be distributed uniformly to cover every leaf to obtain best results. Use of coarse spray nozzles is recommended. Satisfactory defoliation is usually secured with one application; however, with dense cotton a second may be required.

Do not apply within 7 days of harvest. Do not graze treated areas or feed gin waste to livestock.

Wash spray equipment thoroughly after each use. Empty container must be thoroughly washed and decontaminated before reuse. Store in original container. Do not contaminate feed and foodstuffs.

Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty drums. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

SWALLOWED

In case of contact, plenty of water for 15 minutes before reuse.

vegetation. Spray,

WARRANTY OF MERCHANTABILITY, EXPRESSLY EXTEND BEYOND THE LIMITS OF THE STATEMENTS

S.A. BY **WESTERN FARM SERVICE, INC. 2401 Crow Canyon Road, San Ramon, CA. 94583**