PUNCIONE POSSIBLE PROPERTY OF THE PROPERTY OF

HARMFUL IF SWALLOWED

Avoid contact with skin or clothing. Avoid inhalation. In case of contact, flush skin with plenty of water; for eyes, flush with plenty of water for 15 minutes. Get medical attention. Wash contaminated clothing before reuse.

This product is non-selective and is harmful to all vegetation. Spray, drift, seepage and drainage will injure or kill all vegetation.

NOTICE OF WARRANTY

WESTERN FARM SERVICE, INC. MAKES NO WARRANTY OF MER-CHANTABILITY, FITNESS FOR ANY PURPOSE OF OTHERWISE, EXPRESS OR IMPLIED concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.



COTTON DEFOLIANT 2

Contains 2 Pounds Sodium Chlorate per Gallon

ACTIVE INGREDIENT Sodium Chlorate (NaClO ₃)		By Weight 20.2%
INERT INGREDIENTS		79.8%
	TOTAL	100.0%
EPA Reg. No. 11656-45 AA	NET CONTENTS:	GALS.

CAUTION:

KEEP OUT OF REACH OF CHILDREN

Best defoliation of content of the plant are defoliant late to insure percent and preferably time of application. You least 30 days old. The rement Station or State Action

FOR AIRPLANE A
DEFOLIANT in 10-15 ga
TION: Apply 2-2½ gal
water per acre. The defoliation
Satisfactory defoliation
with dense cotton a secon

Do not apply within feed gin waste to livest

Wash spray equipromust be thoroughly was original container. Do n

Do not contaminate wastes. Apply this produced

Do not reuse empty by perforating or crush supplies.

WFS 45-01 8-75

MANUFACTURED IN U.S.A. BY WESTERN FARM SERVICE, INC. 2401 Crow Canyon Road, San Ramon, CA



F SWALLOWED

n. In case of contact, enty of water for 15 othing before reuse.

Il vegetation. Spray, n.

ARRANTY OF MER-IERWISE, EXPRESS extend beyond the with the statements

COTTON DEFOLIANT 2

Contains 2 Pounds Sodium Chlorate per Gallon

ACTIVE INGREDIENT Sodium Chlorate (NaClO ₃)		By Weight 20.2%
INERT INGREDIENTS		79.8%
	TOTAL	100.0%
EPA Reg. No. 11656-45 AA	NET CONTENTS:	GALS.

CAUTION:

KEEP OUT OF REACH OF CHILDREN

WFS 45-01 8-75

DIRECTIONS FOR USE

Best defoliation occurs on cotton when the nitrogen and moisture content of the plant are nearly or completely exhausted. It is better to apply derolant late to insure a higher yield of better quality fiber. At least 75 percent and preferably 85 to 90 percent of the bolls should be open at the time of application. Youngest bolls expected to make cotton should be at least 30 days old. The recommendations of your State Agricultural Experiment Station or State Agriculture Extension Service should be followed.

FOR AIRPLANE APPLICATION: Apply 2-2½ gallons of COTTON DEFOLIANT in 10-15 gallons of water per acre. FOR GROUND APPLICATION: Apply 2-2½ gallons of COTTON DEFOLIANT in 15-30 gallons of water per acre. The defoliant must be distributed uniformly to cover every leaf to obtain best results. Use of coarse spray nozzles is recommended. Satisfactory defoliation is usually secured with one application; however, with dense cotton a second may be required.

Do not apply within 7 days of harvest. Do not graze treated areas or feed gin waste to livestock.

Wash spray equipment thoroughly after each use. Empty container must be thoroughly washed and decontaminated before reuse. Store in original container. Do not contaminate feed and foodstuffs.

Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty drums. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

SA BY WESTERN FARM SERVICE, INC. 2401 Crow Canyon Road, San Ramon, CA. 94583