

11/26/75

MAY CAUSE SKIN IRRITATION HARMFUL IF SWALLOWED

Avoid contact with skin or clothing. Avoid inhalation. In case of contact, flush skin with plenty of water; for eyes, flush with plenty of water for 15 minutes. Get medical attention. Wash contaminated clothing before reuse.

This product is non-selective and is harmful to all vegetation. Spray, drift, seepage and drainage will injure or kill all vegetation.

NOTICE OF WARRANTY

WESTERN FARM SERVICE, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE OF OTHERWISE, EXPRESS OR IMPLIED concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.



COTTON DEFOLIANT 3

Contains 2 Pounds Sodium Chlorate per Gallon

| | |
|---|------------------|
| ACTIVE INGREDIENT | By Weight |
| Sodium Chlorate (NaClO ₃) | 20.1% |
| INERT INGREDIENTS | 79.9% |
| TOTAL | 100.0% |

EPA Reg. No. 11656-46 AA

NET CONTENTS GALS.

CAUTION:
KEEP OUT OF REACH OF CHILDREN

WFS 46-01 8-75

MANUFACTURED IN U.S.A. BY **WESTERN FARM SERVICE, INC. 2401 Crow Canyon Road, San Ramon, CA**

Best defoliation occurs when the leaf content of the plant are not less than 8 percent and preferably 8-10 percent at time of application. Your cotton should be at least 30 days old. The recommended application rate is 2-2 1/2 gallons per acre. For more information contact your County Agent, Extension Station or State Agricultural Experiment Station.

FOR AIRPLANE APPLICATION: Apply 2-2 1/2 gallons of DEFOLIANT in 10-15 gallons of water per acre. The defoliant should be applied to the leaf to obtain best results. Satisfactory defoliation is usually obtained with dense cotton a second year.

Do not apply within 30 days of feed gin waste to livestock.

Wash spray equipment thoroughly with water after use. Do not use original container. Do not reuse.

Do not contaminate water, food, or feed with spray wastes. Apply this product to the ground.

Do not reuse empty container. Do not crush or perforate container. Do not burn.



COTTON DEFOLIANT 3

Contains 2 Pounds Sodium Chlorate per Gallon

| | |
|---|------------------|
| ACTIVE INGREDIENT | By Weight |
| Sodium Chlorate (NaClO ₃) | 20.1% |
| INERT INGREDIENTS | 79.9% |
| TOTAL | 100.0% |

EPA Reg. No. 11656-46 AA

NET CONTENTS GALS.

CAUTION:
KEEP OUT OF REACH OF CHILDREN

WFS 46-01 8-75

DIRECTIONS FOR USE

Best defoliation occurs on cotton when the nitrogen and moisture content of the plant are nearly or completely exhausted. It is better to apply defoliant late to insure a higher yield of better quality fiber. At least 75 percent and preferably 85 to 90 percent of the bolls should be open at the time of application. Youngest bolls expected to make cotton should be at least 30 days old. The recommendations of your State Agricultural Experiment Station or State Agriculture Extension Service should be followed.

FOR AIRPLANE APPLICATION: Apply 2-2½ gallons of COTTON DEFOLIANT in 10-15 gallons of water per acre. **FOR GROUND APPLICATION:** Apply 2-2½ gallons of COTTON DEFOLIANT in 15-30 gallons of water per acre. The defoliant must be distributed uniformly to cover every leaf to obtain best results. Use of coarse spray nozzles is recommended. Satisfactory defoliation is usually secured with one application; however, with dense cotton a second may be required.

Do not apply within 7 days of harvest. Do not graze treated areas or feed gin waste to livestock.

Wash spray equipment thoroughly after each use. Empty container must be thoroughly washed and decontaminated before reuse. Store in original container. Do not contaminate feed and foodstuffs.

Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty drums. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

MADE BY **WESTERN FARM SERVICE, INC. 2401 Crow Canyon Road, San Ramon, CA. 94583**