

DIRECTIONS FOR USE

APPLES: Powdery mildew — 0.5 to 3 gal. per 100 gallons of spray. Apply in prepink, pink and calyx stages and throughout the growing season as needed. Blotch — 2 gal. per 100 gallons spray. Apply in pre-bloom and petal fall stages and ten days later. Leaf spots — 7.5 to 10 gal. per 100 gallons spray in delayed dormant stage or 1/2 gal. per 100 gallons spray in growing season. **Dormant or Delayed Dormant** — Pear psylla, pear psylla eggs, blister mites, rust mites — 4 gal. per 100 gallons plus 1-1/2 gal. superior type oil per 100 gallons. Scab — 6 to 11 gal. per 100 gallons spray. For use in prepink, pink and calyx periods, use 0.5 to 3 gal. per 100 gallons spray. **NOTE:** Use on Delicious apples may result in injury. Apply 1.5 to 2.0 gal. in delayed dormant, prepink and pink stages. Scale insects, aphids — 3 gal. per 100 gallons spray with 1-1/2 gal. superior type oil as conventional full gallonage spray. POLY-SUL may be used alone at 6 to 11 gal. per 100 gallons for control of scale insects, blister mites and rust mites as a full gallonage spray.

BLACKBERRIES: Anthracnose, powdery mildew, rust — 6 to 12 gal. per 100 gallons spray. Apply in dormant or delayed dormant (green bud) stages. Use 0.5 gals. per 100 gallons spray when fruiting canes are about one foot long and before blossoms have opened. Cane blight, overwintering fungus spores — 9.5 to 15.6 quarts per 100 gallons spray. Apply at spring bud swell. Repeat in late fall dormant period.

GRAPES: Anthracnose, powdery mildew — 2 qts. per 100 gallons spray when new shoots are 4 to 6 inches long. Use 1 pt. plus 4 lbs. sulfur per 100 gallons on mature foliage.

NECTARINES: Brown rot, leaf spot, shothole — 7 gal. in 100 gallons spray. Apply 3 to 5 times at weekly intervals before harvest. Leaf curl — 6 gal. in 100 gallons spray. Apply in early winter and late dormant periods. Powdery mildew — 2 to 3 qts. in 100 gallons spray. Apply when disease appears.

PEACHES: Brown rot, leaf spots — 2 qts. per 100 gallons spray. Apply 3 to 5 times at weekly intervals before harvest. Brown rot, blossom blight — 1 pint of POLY-SUL per 8 to 16 gallons spray. Apply in pre-bloom, early bloom and full bloom stages. Leaf curl, scab, shothole — 3.9 to 12 gal. per 100 gallons spray. For leaf curl, use 8 gal. per 100 gallons spray in dormant period. Powdery mildew — 7.5 in 100 gallons spray at delayed dormant stage and 4 to 5 gal. at green tip stage. Use 1.5 qts. on summer foliage. **Dormant or Delayed Dormant Sprays:** Scale insects, silver mites, peach twig borer, aphids — 6 to 11 gal. per 100 gallons spray or 3 gal. per 100 gallons with 1 1/2 gal. superior type oil for additional control of brown and red mites.

WFS 51 01 9 76

WESTERN FARM SERVICE, INC.



POLY-SUL

Fungicide-Insecticide-Miticide

ACTIVE INGREDIENT	BY WEIGHT
Calcium polysulfide	29%
INERT INGREDIENTS	71%
TOTAL	100%

REGISTERED UNDER
EPA REG. NO. 11656-51

Contains calcium and sulfur expressed as gypsum 3.0 lbs. per gallon. Other combined sulfur 1.9 lbs. per gal. Product Weight 10.59 pounds per gallon at 58° F.

EPA Reg. No. 11656-51 NET CONTENTS ___ GALS

Keep Out of Reach of Children

CAUTION

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

CAUTION: Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur is safe for that crop or in that locality. Do not apply oil following POLY-SUL, nor POLY-SUL following oil, in foliage period.

FOR USE AS A SOIL AMENDMENT: Alkaline soil correction and improvement of water penetration: 10 to 30 gallons per acre in irrigation water on growing plants. Make 2 to 3 applications.

For pre-planting treatment: 20 to 30 gallons per acre in irrigation water.

PEARS: Powdery mildew, scab — 6 to 11 gal. per 100 gallons as a full coverage dormant or delayed dormant spray. For powdery mildew, continue spraying throughout growing season as needed. **NOTE:** Use on d'anjou pears after the calyx period may result in injury. **Dormant or Delayed Dormant:** Pear psylla, pear psylla eggs, blister mites, rust mites — 4 gal. per 100 gallons spray with 1-1/2 gal. superior type oil. Scale insects, aphids — 3 gal. per 100 gallons spray with 1-1/2 gal. superior type oil as a conventional full gallonage spray. POLY-SUL may be used alone at 6 to 11 gal. per 100 gallons spray for control of scale insects, scab, blister mites and rust mites as a full gallonage spray. **Post-Harvest Spray** — Pear psylla, rust mite — 4 gal. per 100 gallons spray with 3/4 gal. superior type oil.

PLUMS: Blotch, powdery mildew — 2 qts. per 100 gallons spray. Begin 3 to 5 weeks before harvest. Repeat weekly until picking. Brown rot (fruit rot) — 1 pt. per 16 gallons spray. Apply 20, 10 and 2 days before harvest. Leaf curl — 10 to 12 gal. per 100 gallons spray. Apply in dormant stage. **Dormant or Delayed Dormant Sprays** — Scale insects, silver mites, peach twig borers, aphids — 6 to 11 gal. per 100 gallons spray or 3 gal. with 1 1/2 gal. superior type oil per 100 gallons spray for additional control of brown and red mites.

PRUNES: Black knot — 0.5 pts. per 1 gallon spray. Apply in green tip stage. Brown rot (fruit rot), leaf spot, powdery mildew — 2 qts. per 100 gallons spray. Begin 3 to 5 weeks before harvest. Apply at weekly intervals until picking. **Dormant or Delayed Dormant Sprays** — Scale insects, silver mites, peach twig borer, aphids — 6 to 11 gal. in 100 gallons spray or 3 gal. with 1-1/2 gal. of superior type oil for additional control of brown and red mites.

RASPBERRIES: Anthracnose, powdery mildew — 1 pt. per 1.5 gallons spray. Spray once when new canes are 8 to 12 inches long, or, for anthracnose, use 1 pt. per gallon and apply when buds begin to break. Cane blight, overwintering fungus spores — 1 to 3.75 pints per 12 gallons spray. Apply at spring bud swell. Repeat in late fall dormant period. Spur blight — 10 to 12 gal. per 100 gallons spray. Apply in delayed dormant stage when buds begin to break and show silver.

NOTICE OF WARRANTY

WESTERN FARM SERVICE, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

2401 CROW CANYON ROAD,

SAN RAMON, CA. 94583