



## DIRECTIONS FOR USE

**ALFALFA:** Alfalfa Caterpillar, Powdery Mildew, Rust — Apply 10 to 20 lbs. per acre. Use restricted to Arizona and West Coast only.

**BROCCOLI, CABBAGE, CAULIFLOWER:** Powdery Mildew, Cabbage Looper, Diamond-back Moth Larvae, Imported Cabbage Worm — Apply 30 to 50 lbs. per acre.

**GRAPES:** Grape Leafroller, Omnivorous Leaf Roller, Powdery Mildew, Rust — Apply 20 to 50 lbs. per acre. Start treatment as soon as possible after hatching and before larvae are protected by leaf folds. Do not allow workers to enter treated areas to engage in any activity requiring substantial contact with treated foliage for 1 day following application. Use restricted to California.

**ORANGES:** Fruit Tree Leafroller — Apply 25 to 50 lbs. per acre. Orange Dog — Apply 20 to 50 lbs. per acre. Use restricted to Arizona and California.

**STRAWBERRIES:** Powdery Mildew — Apply 25 to 50 lbs. per acre. Sulfur may injure strawberries under certain climatic conditions.

**TOMATOES:** Tomato Hornworm. Apply 20 to 50 lbs. per acre.

# DIPEL<sup>®</sup> SULFUR

## 150-40 DUST INSECTICIDE FUNGICIDE

ACTIVE INGREDIENTS	By Weight
Bacillus thuringiensis, Berliner, Potency of 320 International Units per mg. (at least 0.5 billion viable spores per g.)*	0.064%
Sulfur	39.200%
INERT INGREDIENTS	60.736%
TOTAL	100.000%

\*Equivalent to potency of 0.15 billion International Units per pound of this product.

EPA REG. NO. 11656-52 AA      NET WEIGHT      LBS.

### CAUTION: KEEP OUT OF REACH OF CHILDREN

**CAUTION:** Avoid inhalation of dust or contact with eyes or open wounds. Avoid prolonged or repeated contact with skin.

Store bag in a cool dry place.

Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. Do not use sulfur on any crop unless local use has proven that sulfur is safe in that locality. Consult your State Agricultural Extension Service or Experiment Station for additional information consistent with label cautions and directions. Do not use within 2 weeks of an oil treatment; on citrus within 3-6 weeks. When grown for processing, consult the processor before applying sulfur. Sulfur may burn foliage when temperature is high. Do not make sulfur applications at such times.

### ENVIRONMENTAL AND DISPOSAL CAUTIONS

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not re-use container. Burn bag when empty. Keep out of smoke.

Dipel is the registered trademark of Abbott Laboratories, North Chicago, Illinois 60064

### NOTICE OF WARRANTY

WESTERN FARM SERVICE MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

## WESTERN FARM SERVICE, INC.

2401 Crow Canyon Road  
San Ramon, California 94583