

WESTERN



ACCEPTED

OCT 6 1987

FARM SERVICE

The Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 11656-53

BTB SULFUR

150-75 DUST INSECTICIDE FUNGICIDE

ACTIVE INGREDIENTS	By Weight
Bacillus thuringiensis, Berliner, Potency of 320 International Units per mg. (at least 0.5 billion viable spores per g.)	0.064%
Sulfur	73.500%
INERT INGREDIENTS	25.436%
TOTAL	100.000%

*Equivalent to potency of 0.15 billion International Units per pound of this product.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FOR AGRICULTURAL USE ONLY OR FOR SALE TO USE AND STORAGE BY SERVICE PERSONS ONLY. DO NOT STORE IN AREAS ACCESSIBLE TO CHILDREN.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

STATEMENT OF PRACTICAL TREATMENT

If swallowed - Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled - Remove victim to fresh air. Apply respiration if indicated.

If on skin - Wash contaminated skin with soap and water.

If in eyes - Flush eyes with plenty of water. Get medical attention immediately.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION!

Avoid inhalation of dust or contact with eyes or open wounds. Avoid prolonged or repeated contact with skin. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS.

REENTRY STATEMENTS

Do not enter treated areas without protective clothing until dusts have settled.

Because certain states may require more restrictive intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE

Keep in original container.
Do not reuse empty container.
Store in cool, dry place.
Protect from excessive heat.

For help with any spill, leak, fire or exposure involving this material, call day or night 800-424-9300.

DISPOSAL

Wastes resulting from the use of this product may be disposed of at an approved waste disposal facility.

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, if allowed by State and local authorities. If burned, stay out of smoke.

BEST AVAILABLE COPY

CAUTIONS IN USING:

Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. Do not use sulfur on any crop unless it has been proven that sulfur is safe in that crop. Consult your State Agricultural Extension or Experiment Station for additional information consistent with the label cautions and directions. When grown for processing, consult the processor before applying sulfur. Sulfur injury to foliage when temperature is high. Do not apply at such times.

GENERAL USE DIRECTIONS

GRAPES: For Powdery Mildew apply 30 lbs. per acre when shoots are 6 inches tall. Repeat before blossoms open and continue at 2 week intervals as necessary.

For Powdery Mildew and Grape Leafhopper apply 30 lbs. per acre as possible after egg hatch and before leaves are protected by leaf folds. Do not allow treated areas within 1 day (24 hours) of application to engage in any activity which involves substantial contact with treated foliage. NOTE: Concord and other Vitis species may be injured.

MELONS: For Powdery Mildew apply 30 lbs. per acre. For Powdery Mildew and Citrus Red Mite apply 30 lbs. per acre. Begin when leaves appear and repeat at weekly intervals. NOT use within 2 weeks of an oil spray.

NOTE: Certain varieties of Melons are susceptible to Sulfur damage.

NOTICE OF WARRANTY

WESTERN FARM SERVICE MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR A PARTICULAR USE OR OTHERWISE, EXPRESS OR IMPLIED. THE USER'S USE OF THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS OF USE ACCORD WITH THE STATEMENTS MADE ON THE LABEL.

WESTERN FARM SERVICE,
3075 Citrus Circle, Suite 100
Walnut Creek, California 94694

EPA Est. No. 11656-53
EPA Est. No. 11656-CA-0

NET WEIGHT: _____ POUNDS