

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION!

May irritate eyes, nose, throat and skin. Harmful if inhaled or swallowed. Do not inhale dust or spray. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. The consumption of alcoholic beverages increases the possibility of harm.

ENVIRONMENTAL HAZARDS

This pesticide product is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. Drift or runoffs from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water or wetlands.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS.

REENTRY STATEMENT

Do not reenter treated areas within 24 hours following application unless protective clothing is worn.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Use Precautions

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not rotate treated crops with crops other than those with registered Thiram uses.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food for feed by storage or disposal.

STORAGE


Keep pesticide in original container. Do not reuse container. Store in cool, dry place. Protect from excessive heat.

For help with any spill, leak, fire or exposure involving this material, call day or night
800-424-9300.

DISPOSAL

Do not dispose of product or container in or near any body of water. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.



THIRAM 65%
Wettable Powder Fungicide

ACTIVE INGREDIENT:	BY WEIGHT
Thiram (Tetramethylthiuram-disulfide).....	65%
INERT INGREDIENTS.....	35%
TOTAL	100%

KEEP OUT OF REACH OF CHILDREN
CAUTION
FOR AGRICULTURAL USE ONLY OR FOR SALE TO USE, AND STORAGE BY SERVICE PERSONS ONLY. DO NOT STORE IN AREAS ACCESSIBLE TO CHILDREN.
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Harmful if swallowed. Wash thoroughly after handling.

IF INHALED: Harmful if inhaled. Avoid breathing dust. Remove contaminated clothing and wash before reuse.

IF ON SKIN: Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling.

IF IN EYES: Causes eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

BEST AVAILABLE COPY

WESTERN FARM SERVICE, INC.

3075 Citrus Circle
Walnut Creek, CA 94598

EPA REG. NO. 11656-68

NET WEIGHT POUNDS

THIRAM 65%
Wettable Powder Fungicide
DIRECTIONS FOR USE

Thiram may be washed into the spray tank through a deep screen or be made into a thick paste by mixing with a little water and slaking, then pouring into the spray tank while filling. Addition of a Spreader-Sticker to the spray tank will promote uniform coverage on foliage and fruit. Use Thiram at 1½ to 2 lbs. per 100 gals. of water, unless otherwise stated, employing higher dosage when disease conditions are severe and reducing to lower dosage when situation is normal for disease development. Observe all cautions and limitations on labeling of all products used in mixtures.

APPLES: Scab—Apply as protective fungicide in pre-bloom, calyx and early cover sprays. Use 1 to 1½ lbs. in later cover sprays.

As a dust (In New England States only)—Use a mixture containing 7.5% to 10% Thiram (4.88% to 6.5% active) mixed with a suitable diluent such as talc or pyrophyllite at 35 to 40 lbs. per acre. Good coverage of foliage and developing fruit is required for best results. (Timing same as for sprays.)

Cedar Apple Rust—Apply in pre-bloom, calyx and first cover sprays. If recommended by local authorities, apply a special bloom spray at a reduced rate of ¼ lb. per 100 gals. of water. Keep foliage and fruit covered during infection periods.

Rates of application of dusts (New England States only) for rust control are the same as for scab, and time is same as given above for sprays.

Other Apple Diseases—For control of Black Rot (fruit and foliage), Fly Speck, Sooty Blotch, Bitter Rot, Brooks Spot, Apple Blotch, Botryosphaeria Spot, and Black Pox—Make spray applications as indicated for scab control since these are effective in controlling the other diseases listed. Whenever weather conditions favor disease, shorten interval between applications. Thiram is a protective fungicide; hence, applications are most effective when made in advance of infection periods.

PEACHES: Brown Rot—For blossom blight, apply at 3- to 4-day intervals during bloom, timed to keep newly opened blossoms covered. For fruit rot, apply at petal-fall and two weeks later. Make one or more additional applications at 7-day intervals beginning about one month before harvest, or earlier if rot is apparent on green fruit. **Rhizopus Rot**—Apply as directed for fruit phase of brown rot. **Scab**—Apply at petal-fall and repeat 3 to 6 times at 10- to 14-day intervals or until disease is well under control. Do not apply later than 7 days before harvest.

TOMATOES (Florida only) For control of gray mold (Botrytis) and Rhizoctonia fruit rot, Thiram is a supplement to the regular spray program employed for control of gray leaf spot (Stemphylium), late blight, and other diseases.

Use Thiram alone for gray mold and Rhizoctonia at the rate of 1½ lbs. per 100 gals. of water. If gray leaf spot, late blight, or other foliage diseases are active, use combination spray of 1 lb. Thiram plus 1½ lbs. of MANZATE[®] or MANZATE[®] D Maneb Fungicides or MANZATE[®] 200 Fungicide. Apply at 3- to 5-day intervals, beginning when plants are set in the field and continuing as long as disease threatens. Use sufficient spray to cover all portions of plants thoroughly. If use in combination spray with MANZATE[®] products, do not apply within 5 days of harvest.

CELERY: For control of early blight (Cercospora), late blight (Septoria) and Rhizoctonia, use 1½ lbs. Thiram in 100 gals. of water. **Plant Beds**—Apply 15 to 25 gals. per 1200 sq. ft. of bed at 3-day intervals; shorten interval under conditions of severe disease incidence. **Field (Florida only)**—Continue treatments after plants are set in the field, applying as a full coverage spray (up to 200 gals. per acre) as frequently as needed (weekly under average conditions; every 3 to 5 days under severe conditions). Do not apply later than 7 days before harvest. Remove residues by stripping, trimming and washing.

Plant beds that are treated with this product must bear distinct labeling that instructs:

CAUTION: Treated with thiram. Wear rubber gloves and protective clothing when handling celery treated with thiram. Direct exposure to skin may result in severe dermatitis.

BEST AVAILABLE COPY

NOTICE OF WARRANTY

WESTERN FARM SERVICE MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORD WITH THE STATEMENTS MADE ON THIS LABEL.



SUPPLEMENTAL LABEL
THIRAM 65% WETTABLE POWDER
FUNGICIDE

FOR CONTROL OF GRAY MOLD
ON STRAWBERRIES
IN CALIFORNIA ONLY

Thiram Fungicide is recommended for use on strawberries for the control of gray mold (Botrytis).

AS A SPRAY - Use 2 to 2½ lbs. Thiram per 100 gals. of water;
apply at rates up to 200 gals. per acre.

WHEN TO APPLY: Begin application at early blossom stage.
Repeat at 10 day intervals as needed. Good coverage of
strawberry buds, blossoms and developing fruit is required
for best results.

Do not apply within 3 days of picking.

IMPORTANT

BEFORE USING THIS PRODUCT READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS
AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product
which do not appear on package label. Follow the instructions carefully.

EPA Reg. No. 11656-68

WESTERN FARM SERVICE, INC.
3075 Citrus Circle, Suite 195
Walnut Creek, California 94598

BEST AVAILABLE COPY



THIRAM 65% WETTABLE POWDER FUNGICIDE

SUPPLEMENTAL LABELING
EPA REG. NO. 11656-68

FOR CONTROL OF
CERTAIN DISEASES OF BANANAS.

Thiram Fungicide is recommended for pre-harvest use on bananas for the control of fruit spots caused by organisms such as Pyricularia grisea and Fursarium roseum.

HOW TO USE

Use 1 lb. Thiram in 5 gals. of water. Apply with a knapsack sprayer at weekly intervals, beginning when the banana flower emerges. Direct spray to area of developing fruit. Do not make more than 3 applications.

IMPORTANT

BEFORE USING THIRAM, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.



SUPPLEMENTAL LABEL

THIRAM 65% WETTABLE POWDER
FUNGICIDE

FOR CONTROL OF
GRAY MOLD ON STRAWBERRIES

Thiram Fungicide is recommended for use on strawberries for the control of gray mold (Botrytis).

HOW TO USE

As a Spray - Use 2 to 2½ lbs. Thiram per 100 gals. of water; apply at rates up to 200 gals. per acre.

As a Dust - Use a prepared dust containing 7.5% to 10% Thiram (4.88% to 6.5% active) mixed with a suitable diluent such as talc or pyrophyllite, at the rate of 40 lbs. per acre.

WHEN TO APPLY

Make 3 to 5 applications at 10-day intervals beginning at early blossom stage. Good coverage of strawberry buds, blossoms and developing fruit is required for best results.

If applied within 3 days of picking, remove residues by washing.

IMPORTANT

BEFORE USING THIS PRODUCT READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on package label. Follow the instructions carefully.

EPA Reg. No. 11656-68

WESTERN FARM SERVICE, INC.
3075 Citrus Circle, Suite 195
Walnut Creek, California 94598