



WFSI SEVIN*-SULFUR 10-50 DUST

ACTIVE INGREDIENTS:

*Carbaryl (1-naphthyl-N-methyl carbamate).....	10%
Sulfur.....	50%
INERT INGREDIENTS.....	40%

TOTAL 100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

FOR AGRICULTURAL USE ONLY OR FOR SALE TO, USE, AND STORAGE BY SERVICE PERSONS ONLY. DO NOT STORE IN AREAS ACCESSIBLE TO CHILDREN.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals

CAUTION!

May be harmful if swallowed, inhaled or absorbed through skin. Do not take internally. Do not breathe dust. Skin contact may be harmful. Avoid unnecessary contact. Wash hands before eating or smoking. Take a shower or bath after work. Wear regular long sleeved work clothing. Change clothing daily. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Do not apply during periods of bloom. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS

REENTRY STATEMENTS

Do not enter treated area without protective clothing until dusts have settled.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE

Keep in original container. Do not reuse empty container. Store in cool, dry place. Protect from excessive heat. For help with any spill, leak, fire or exposure involving this material, call day or night 800-424-9206.

DISPOSAL

Wastes resulting from the use of this product may be disposed of at an approved waste disposal facility.

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

GENERAL USE DIRECTIONS

May be applied by ground or air equipment. This product may be applied up to a maximum permissible rate per acre, expressed as (#) after each crop class.

Subsequent planting of feed and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

Do not use on cucurbits or other plants susceptible to sulfur injury.

BEAN: Mexican bean beetle, leafhoppers, powdery mildew - 10 lbs. per acre. Lygus, stink bugs - 20 lbs. per acre. Apply thoroughly at first signs of disease or insect infestation. When necessary, repeat every 7 to 10 days. No time limitation. (20#)

COTTON (early season): Thrips, cotton fleahopper - 15 to 20 lbs. per acre as soon as insects appear. (Mid and late season): Boll weevil, bollworm, pink bollworm,

Cotton leafworm, lygus bug, cotton leaf perforator, aphid - 20 to 25 lbs. per acre. Do not apply within 7 days of harvest.

CALIFORNIA ONLY: Bollworm - 40 lbs. per acre. Apply when insects appear and, if necessary, repeat at 5 to 7 day intervals. Aphid population will usually be suppressed when this material is used on a regular schedule of treatments. Do not apply within 7 days of harvest. (25#)

GENERAL USE DIRECTIONS (Cont'd.)

GRAPES: Grape leaf folder, Grape leafhopper, Grape leaf skeletonizer, red-banded leaf roller, June beetles, Japanese beetle, grape berry moth, powdery mildew - 10 to 30 lbs. per acre. Thorough coverage is necessary. Workers are not to enter treated area to engage in any activity requiring substantial contact with treated foliage until 2 days after application. (30#)

PEACHES, NECTARINES: Oriental fruit moth, Peach twig borer, Plum curculio, catfacing insects, periodical cicada, powdery mildew, brown rot - 35 to 60 lbs. per acre. Apply thoroughly at first signs of disease or insect infestation. When necessary, repeat at 7 to 10 day intervals. Workers are not to enter treated area to engage in any activity requiring substantial contact with treated foliage until 2 days after application. Do not apply within 1 day of harvest for peaches, or within 3 days of harvest for nectarines. (60#)

PLUMS, PRUNES: Fruit tree leaf roller, Peach twig borer, plum curculio, red-banded leaf roller, codling moth, Eastern tent caterpillar, Orange tortrix, Western tussock moth - 40 to 60 lbs. per acre. Apply when insects appear and, when necessary, repeat at 7 to 10 day intervals. Make full coverage application. Do not apply within 1 day of harvest. (60#)

POTATOES: Colorado potato beetle - 10 lbs. per acre. Leafhoppers, flea beetle, lygus bugs - 20 lbs. per acre. Apply when insects appear and, if needed, repeat application at 7 to 10 day intervals. Apply thoroughly to upper and lower leaf surfaces. (20#)

TOMATOES: Colorado potato beetle - 10 lbs. per acre. Flea beetles, leafhoppers, cutworms - 20 lbs. per acre. Tomato fruitworm, fall armyworm, tomato hornworm, European corn borer, stink bugs, lace bugs, tarnished plant bugs - 30 to 40 lbs. per acre. Because of the sulfur content of this dust, check with your canner of fertilizer before applying to tomatoes. No time limitation. (40#)

*SEVIN is a registered trademark of Union Carbide Corporation.

NOTICE OF WARRANTY

WESTERN FARM SERVICE MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR CONTAINED, EXPRESS OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORD WITH THE STATEMENTS MADE ON THIS LABEL.

NET CONTENTS: _____ POUNDS

WESTERN FARM SERVICE, INC.
3075 Citrus Circle, Suite 195
Walnut Creek, CA 94598

BEST AVAILABLE COPY



WFSI SEVIN*-SULFUR 10-50 DUST

ACTIVE INGREDIENTS:	BY WT.
*Carbaryl (1-naphthyl-N-methyl carbamate).....	10%
Sulfur.....	50%
INERT INGREDIENTS.....	40%
TOTAL	100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

FOR AGRICULTURAL USE ONLY OR FOR SALE TO, USE, AND STORAGE BY SERVICE PERSONS ONLY. DO NOT STORE IN AREAS ACCESSIBLE TO CHILDREN.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals

CAUTION!

May be harmful if swallowed, inhaled or absorbed through skin. Do not take internally. Do not breathe dust. Skin contact may be harmful. Avoid unnecessary contact. Wash hands before eating or smoking. Take a shower or bath after work. Wear regular long sleeved work clothing. Change clothing daily. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Do not apply during periods of bloom. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS

REENTRY STATEMENTS

Do not enter treated area without protective clothing until dusts have settled.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE

Keep in original container. Do not reuse empty container. Store in cool, dry place. Protect from excessive heat. For help with any spill, leak, fire or exposure involving this material, call day or night 800-424-9366.

DISPOSAL

Wastes resulting from the use of this product may be disposed of at an approved waste disposal facility.

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

GENERAL USE DIRECTIONS

May be applied by ground or air equipment. This product may be applied up to a maximum permissible rate per acre, expressed as (f) after each crop class:

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

Do not use on cucurbits or other plants susceptible to sulfur injury.

BEANS: Mexican bean beetle, leafhoppers, powdery mildew - 10 lbs. per acre. Lygus, stink bugs - 20 lbs. per acre. Apply thoroughly at first signs of disease or insect infestation. When necessary, repeat every 7 to 10 days. No time limitation. (20r)

COTTON (Early season): Thrips, cotton flea-hopper - 15 to 20 lbs. per acre as soon as insects appear. (Mid and late season): Boll weevil, bollworm, pink bollworm, Cotton leafworm, lygus bug, cotton leaf perforator, aphid - 20 to 25 lbs. per acre. Do not apply within 7 days of harvest.

CALIFORNIA ONLY: Bollworm - 40 lbs. per acre. Apply when insects appear and, if necessary, repeat at 5 to 7 day intervals. Aphid population will usually be suppressed when this material is used on a regular schedule of treatments. Do not apply within 7 days of harvest. (25a)

GENERAL USE DIRECTIONS (Cont'd.)

GRAPES: Grape leaf folder, Grape leafhopper, Grape leaf skeletonizer, red-banded leaf roller, June beetles, Japanese beetle, grape berry moth, powdery mildew - 10 to 30 lbs. per acre. Thorough coverage is necessary. Workers are not to enter treated area to engage in any activity requiring substantial contact with treated foliage until 2 days after application. (30r)

PEACHES, NECTARINES: Oriental fruit moth, Peach twig borer, Plum curculio, catfacing insects, periodical cicada, powdery mildew, brown rot - 25 to 60 lbs. per acre. Apply thoroughly at first signs of disease or insect infestation. When necessary, repeat at 7 to 10 day intervals. Workers are not to enter treated area to engage in any activity requiring substantial contact with treated foliage until 2 days after application. Do not apply within 1 day of harvest for peaches, or within 3 days of harvest for nectarines. (60r)

PLUMS, PRUNES: Fruit tree leaf roller, Peach twig borer, plum curculio, red-banded leaf roller, rolling moth, Eastern tent caterpillar, Orange tortrix, Western tussock moth - 30 to 60 lbs. per acre. Apply when insects appear and, when necessary, repeat at 7 to 10 day intervals. Make full coverage application. Do not apply within 1 day of harvest. (60r)

POTATOES: Colorado potato beetle - 10 lbs. per acre. Leafhopper, flea beetle, lygus bug - 20 lbs. per acre. Apply when insects appear and, if needed, repeat application at 7 to 10 day intervals. Apply thoroughly to upper and lower leaf surfaces. (20r)

TOMATOES: Colorado potato beetle - 10 lbs. per acre. Flea beetles, leafhoppers, cutworms - 20 lbs. per acre. Tomato fruitworm, fall armyworm, tomato hornworm, European corn borer, stink bugs, lace bugs, tarnished plant bugs - 30 to 40 lbs. per acre. Because of the sulfur content of this dust, check with your canner of sprayer before applying to tomatoes. No time limitation. (40r)

*SEVIN is a registered trademark of Union Carbide Corporation.

NOTICE OF WARRANTY

WESTERN FARM SERVICE MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

NET CONTENTS: _____ POUNDS