



WFSI SEVIN* 5 DUST

ACTIVE INGREDIENTS:	BY WT.
Carbaryl (1-naphthyl-N-methyl carbamate).....	5.0%
INERT INGREDIENTS.....	95.0%
TOTAL	100.0%

Under the
 provisions of
 the Federal
 Insecticide Act,
 this product is
 registered for use
 on the following
 crops:

KEEP OUT OF REACH OF CHILDREN

CAUTION

FOR AGRICULTURAL USE ONLY OR FOR SALE TO, USE, AND STORAGE BY SERVICE PERSONS ONLY. DO NOT STORE IN AREAS ACCESSIBLE TO CHILDREN.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

PRECAUTIONARY STATEMENTS
 Hazards to Humans & Domestic Animals

CAUTION!
 May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling and before eating or smoking. Do not take internally. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS
 This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands, except for rice uses. Discharge from rice fields may kill aquatic and estuarine invertebrates. Do not contaminate water by cleaning of equipment or disposal of wastes. Avoid contamination of food, feed, feeding troughs and watering receptacles. Do not apply under possible drift conditions to food, forage or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Protective information may be obtained from your Cooperative Agricultural Extension Service.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE
 Keep in original container. Do not reuse empty container. Store in cool, dry place. Protect from excessive heat. For help with any spill, leak, fire or exposure involving this material, call day or night 800-424-9300.

DISPOSAL
 Wastes resulting from the use of this product may be disposed of at an approved waste disposal facility. Completely empty bag into application equipment. Then dispose of empty bag in sanitary landfill or by incineration, or, if allowed by State and local authorities, burning. If burned, stay out of smoke.

GENERAL USE DIRECTIONS

May be applied by ground or air equipment. This product may be applied up to the maximum permissible rate, per acre, expressed as (4) after each crop claim.

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels).

ALFALFA, CLOVER, COPEAS, PEANUTS, SOYBEANS: Blister beetles, Mexican bean beetle - 10 to 20 lbs. per acre. Alfalfa caterpillar, bean leaf beetle, cucumber beetles, green cloverworm, Japanese beetle, leafhoppers, three-cornered alfalfa hopper, thrips, velvet bean caterpillar - 20 lbs. per acre. Armyworms, corn earworm, stink bugs, webworms - 20 to 30 lbs. per acre. May be applied up to harvest. Some leaf injury may occur if applications are made when tender foliage is wet. For alfalfa and copeas do not apply within 3 days of harvest. No time limitations for clover, peanuts or soybeans. (30)

APPLES: Codling moth, orange tortrix (skin worm) 60 lbs. per acre. Do not use within 1 day of harvest. (80)

APRICOTS: Codling moth, peach twig borer, leaf roller, Diabrotic beetle - 75 to 80 lbs. per acre. Do not apply within 3 days of harvest. (80)

BEANS: Mexican bean beetle - 10 lbs. per acre. Leafhoppers, bean leaf beetle, cucumber beetles, Western bean cutworm, velvet bean caterpillar, flea beetles, Japanese beetle - 20 lbs. per acre. Corn earworm, armyworms, tarnished plant bug, stink bugs - 20 to 30 lbs. per acre. In California for corn earworm, lima bean pod borer, lygus, stink bugs - 40 lbs. per acre. May be applied to harvest. (40)

BLUEBERRIES: Blueberry maggot, cherry and cranberry fruitworms, Japanese beetles - 30 to 40 lbs. per acre. Apply 3 weeks before harvest and repeat 10 days later or as necessary. May be applied up to harvest. (40)

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, EDIBLES: Flea beetles, harlequin bug - 10 to 20 lbs. per acre. Armyworms, corn earworm, imported cabbageworm, young cabbage looper, salt marsh caterpillar - 20 to 40 lbs. per acre. Do not apply within 3 days of harvest. (40)

GENERAL USE DIRECTIONS (Cont'd.)

CORN: Corn earworm, corn rootworm adults, European corn borer, fall armyworm, sap beetles - 30 to 40 lbs. per acre. For insects attacking silks and ears apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. May be applied up to harvest. (40)

CUCUMBERS, MELONS, PUMPKINS, SQUASH: Pickleworm, melonworm - 10 to 25 lbs. per acre. Cucumber beetles, flea beetles, leafhoppers, squash bug - 20 lbs. per acre. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. May be used up to harvest. (20)

GRAPES: Grape leaf folder, grape leafhopper, grape leaf skeletonizer - 20 to 40 lbs. per acre. Apply just before first brood leaf folder larvae emerge from rolls and as needed for leafhoppers. Grape berry moth, Japanese beetle, June beetles, reo-banded leaf roller - 40 lbs. per acre. (60)

LETTUCE, CARROTS: Flea beetles, harlequin bug, leafhoppers - 10 to 20 lbs. per acre. Six spotted leafhopper - 20 to 30 lbs. per acre in a 5 to 7 day schedule. May be used up to harvest on carrots. Do not use on head lettuce within 3 days of harvest or on leaf lettuce within 14 days of harvest. (40)

ORCHARD: Corn earworm, stink bugs - 20 to 40 lbs. per acre. Repeat at 5 to 7 day intervals or as necessary. No time limitation. (40)

PEACHES: Oriental fruit moth, peach twig borer, plum curculionid, catfacing insects, periodical cicada - 70 to 80 lbs. per acre. Do not apply within 1 day before harvest. (80)

POTATO, EGGPLANT, PEPPER: Colorado potato beetle, flea beetles, leafhoppers - 10 to 20 lbs. per acre. Tomato fruit worm, fall armyworm, tomato hornworm, European corn borer, lygus, stink bugs, lace bugs, tarnished plant bugs - 20 to 40 lbs. per acre. (40)

RICE, FORAGE GRASSES, PASTURE: Armyworms, stink bugs, thrips - 20 to 30 lbs. per acre. For optimum thrip control in grasses grown for seed, direct dust into boot. Do not apply to rice within 14 days of harvest. May be applied up to harvest on pasture for forage grasses. Do not use on rice fields which crayfish and/or catfish farming is included in the cultural practice. (30)

SORGHUMS (Including Milo, Grain Sorghum): Armyworms, corn earworm, stink bugs, webworms - 20 to 40 lbs. per acre. Sorghum midge - 30 lbs. per acre. Dust into the forming heads. Treat sorghum twice 3 to 4 days after heads emerge from boot. Do not apply to grain in a grain within 21 days of harvest. No time limitation for forage. (40)

STRAWBERRIES: Meadow Loititebug, strawberry leaf roller, strawberry weevil - 20 to 40 lbs. per acre. Do not apply within 1 day of harvest. (40)

SUGAR BEETS: Armyworms, leafhoppers, beetworms - 20 to 40 lbs. per acre. Do not apply within 14 days of harvest.

TOMATOES: Flea beetle, leafhoppers, tomato fruitworm, fall armyworm, tomato hornworm, European corn borer - 50 to 60 lbs. per acre. No time limitation. (80)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS.

REENTRY STATEMENT

Do not enter treated areas without protective clothing until dusts have settled.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product.

Calif. Reg. No. 11656-50054-AA
 EPA Est. No. 11656-CA-02

NET CONTENTS: _____ POUNDS

WESTERN FARM SERVICE, INC.
 3075 Citrus Circle, Suite 195
 Walnut Creek, CA 94598

NOTICE OF WARRANTY

WESTERN FARM SERVICE MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.