

USE DIRECTIONS

Swimming Pools

To prevent development of troublesome algae such as *Chlorella pyrenoidosa* (green) and *Phormidium inundatum* (black) use Algaecide-Fungicide according to the following directions:

INITIAL DOSE - 2 quarts for each 50,000 gallons of water.

MAINTENANCE DOSE - ½ quart for each 50,000 gallons of water.

Initial dose is used when filling pool at beginning of swimming season. If pool has visible growth, treat with initial dose. Vacuum pool after 24 hours to remove algae debris. If visible algae are still present or reappear, repeat this treatment procedure as necessary. When pool is free of visible algae, use the recommended maintenance dose.

Maintenance dose should be added at 3-5 day intervals. If high temperatures prevail, or pool has unusually heavy use, add maintenance dose more frequently.

If make-up water is added at the rate of more than 10% per week of pool capacity, add ½ quart of Algaecide-Fungicide for each 10,000 gallons of fresh make-up water added per week.

When swimming season is over, add 2 quarts of Algaecide-Fungicide to 50,000 gallons of water in treating water left in pool. This dose helps provide a measure of control of algae growth during the winter months. This treatment will also help the servicing of the pool before it is returned to use the following season.

Rinse empty container thoroughly with water and discard.

First Aid

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly, a large quantity of milk, egg whites, gelatin solution or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

Note to Physician

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

E.P.A. REG. NO.

Instasan

BRAND

ALGA

Active Ingredients:

- p-Alkyl (60% C₁₄, 3 dimethyl benz
- o-Alkyl (68% C₁₂, 3 ethylbenzyl an

INERT INGREDIENTS

DANGER

KEI
stat

INSTASAN® LABORA

CONTENTS: 1 GALLON

ACCEPTED

MAR 28 1972

E.P.A. REG. NO. 11662-5

PROTECT FROM FREEZING

Instasan 20 Algaecide-Fungicide

BRAND CONCENTRATED

ALGAECIDE-FUNGICIDE

Active Ingredients:

n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈) dimethyl benzyl ammonium chlorides	10%
n-Alkyl (68% C ₁₂ , 32% C ₁₄) dimethyl ethylbenzyl ammonium chlorides	10%
INERT INGREDIENTS	80%
	<hr/> 100%

DANGER KEEP OUT OF REACH OF CHILDREN. See other precautionary statements on Right Side Panel.

INSTASAN[®] LABORATORIES DIV. • ENVIRACHEM, INC. • DETROIT, MICH. 48227

CONTENTS: 1 GALLON

PRINTED IN U.S.A.

Industrial

Our Algaecide-Fungicide is the latest development of (black algae), Ph

Initial dose: 1-2 gallons per system. Use this

Periodic Dose: 1 gallon of Fungicide at the

1-3 quarts of Algaecide-Fungicide 7 days or sooner

For best results, do not be alarmed with operation. This is treated and discolored surfaces as living screens and pipes

Fungicide

Algaecide-Fungicide Strain #640 will remove floors, foot basins 1 gallon of water

DANGER

Keep Out Of Reach of eyes, on skin or clothing. Harmful or fatal

This product is toxic to lakes, streams, ponds

ACCEPTED