

SIM-CHEM

ACCEPTED
 1-19-72
 UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC PEST CONTROL REGISTERED UNDER NO. 11682-1

POLYRAM 7 POTATO SEED TREATER

DIRECTIONS FOR USE

POTATO-CUT SEED: Fusarium Seed Piece Decay, Black Leg, Bacterial Common Rot - Apply 1 to 1½ pounds of Seed Treater to each 100 pounds of seed pieces. For effective control, Seed Treater must be applied to entire surface of each piece immediately after cutting. If planting is delayed, follow the recommended storage practices of cut seed pieces in your area.

POTATO-WHOLE SEED: Seed-borne Common Rot - Apply 1 to 1½ pounds of Seed Treater to each 100 pounds of seed. Seed Treater should be applied evenly to entire surface. If the treated whole seed is cut, a second application will be necessary, if Black Leg and Fusarium Seed Piece Decay are to be controlled.

AN AGRICULTURAL FERTILIZER

ACTIVE INGREDIENTS

A mixture of 5.2 parts by weight (83.0%) of ammoniates of [ethylenebis (dithiocarbamate)] zinc with 1 part by weight (16.1%) ethylenebis (dithiocarbamate acid) bimolecular and tri-molecular cyclic anhydrosulfides and disulfides 7.0%

INERT INGREDIENTS 93.0%

Zinc expressed as metallic 1.22%

U.S. Patent Pending
 Contains Polyram® Registered Trade Mark of Basiglio Anilin & Soda Fabrik A.G.

E.P.A. Reg. No. 11682-1

CAUTION

Do not allow this material to drift onto neighboring crops or use in a manner or at a time other than in accordance with directions, because plant injury, excessive residues or other undesirable results may occur.

May be harmful if swallowed or inhaled. May cause irritation of eyes, nose, throat and skin. In case of contact, flush with plenty of water; for eyes get medical attention. Wear rubber gloves and respirator when mixing Seed Treater with potato seed pieces.

Do not reuse container. Destroy when empty.


Do not use treated seed for feed or feed purposes.

CAUTION Keep Out of Reach of Children
HARMFUL IF TAKEN INTERNALLY
 AVOID INHALING DUST . . . AVOID CONTACT WITH SKIN
 AVOID CONTAMINATION OF FEED AND FOOD
 STORE IN A COOL, DRY PLACE - PROTECT FROM EXCESSIVE HEAT

A PRODUCT OF
SIM-CHEM
 MINERALS AND CHEMICALS DIVISION
 A S. S. DUPONT COMPANY
 Post Office Box 870
 MOUNTAIN HOME, IDAHO 83647

NET WEIGHT 20 LBS.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. But seller disclaims any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, as to the use of the product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.



SIM-CHEM

ACCEPTED
1-19-72
11682-1

POLYRAM 7 POTATO SEED TREATER

DIRECTIONS FOR USE

POTATO-CUT SEED: Fusarium Seed Piece Decay, Black Leg, Seed-borne Common Scab - Apply 1 to 1½ pounds of Seed Treater to each 100 pounds of seed pieces. For effective control, Seed Treater must be applied to entire surface of each piece immediately after cutting. If planting is delayed, follow the recommended storage practices of cut seed pieces in your area.

POTATO-WHOLE SEED: Seed-borne Common Scab - Apply 1 to 1½ pounds of Seed Treater to each 100 pounds of seed. Seed Treater should be applied evenly to entire surface. If the treated whole seed is cut, a second application will be necessary, if Black Leg and Fusarium Seed Piece Decay are to be controlled.

AN AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENTS

A mixture of 5.2 parts by weight (83.9%) of ammoniates of [ethylenebis (dithiocarbamate)] zinc with 1 part by weight (16.1%) ethylenebis [dithiocarbamic acid] bimolecular and trimolecular cyclic anhydrosulfides and disulfides 7.0%

INERT INGREDIENTS 93.0%

Zinc expressed as metallic 1.22%

U.S. Patent Pending
Contains Polyram® Registered Trade Mark of Badische Anilin & Soda Fabrik A.G

E.P.A. Reg. No. 11682-1

CAUTION

Do not allow this material to drift onto neighboring crops or use in a manner or at a time other than in accordance with directions, because plant injury, excessive residues or other undesirable results may occur.

May be harmful if swallowed or inhaled. May cause irritation of eyes, nose, throat and skin! In case of contact, flush with plenty of water; for eyes get medical attention. Wear rubber gloves and respirator when mixing Seed Treater with potato seed pieces.

Do not reuse container. Destroy when empty.

Do not use treated seed for food or feed purposes.

CAUTION Keep Out of Reach of Children
HARMFUL IF TAKEN INTERNALLY
AVOID INHALING DUST . . . AVOID CONTACT WITH SKIN
AVOID CONTAMINATION OF FEED AND FOOD
STORE IN A COOL DRY PLACE - PROTECT FROM EXCESSIVE HEAT

A PRODUCT OF
SIM-CHEM
MINERALS AND CHEMICALS DIVISION
J. R. SIMPLOT COMPANY
Post Office Box 810
MOUNTAIN HOME, IDAHO 83647

NET WEIGHT 50 LBS.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.