

SIM-CHEM

CAPTAN 10 DUST POTATO SEED TREATER

ACCEPTED
JAN 10 1973
UNIVERSITY OF IDAHO
FUNGICIDE REGISTRATION DIVISION
FOR REGISTRATION OF CAPTAN 10 DUST
EXAMINATION NO. 14882 SUBJECT
TO ATTACHED COMMENTS

CAUTION

Do not allow this material to drift onto neighboring crops or use in a manner or at a time other than in accordance with directions, because plant injury, excessive residues or other undesirable results may occur.

Do not use treated seed for food or feed purposes.

May be harmful if swallowed or inhaled. May cause irritation of eyes, nose, throat and skin! In case of contact, flush with plenty of water; for eyes get medical attention. Wear rubber gloves and respirator when mixing Seed Treater with potato seed pieces.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Dispose of excess treated seeds or wastes by burial.

Do not reuse container. Destroy when empty.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

AN AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENT:

Captan (N- [(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide) 10%

INERT INGREDIENTS 90%

Not licensed for use in countries where valid patents exist. 100%

E.P.A. Reg. No. 11682-5
SUB REG. NO. 476-655-15154

CAUTION Keep Out of Reach of Children

HARMFUL IF TAKEN INTERNALLY

AVOID INHALING DUST

AVOID CONTACT WITH SKIN

AVOID CONTAMINATION OF FEED AND FOOD

STORE IN A COOL DRY PLACE; PROTECT FROM EXCESSIVE HEAT

A PRODUCT OF
SIM-CHEM

AGRICULTURAL AND CHEMICALS DIVISION • J. R. SIMPLOT COMPANY
Post Office Box 810
Mountain Home, Idaho 83647

NET WEIGHT 50 LBS.

RECOMMENDATIONS


POTATO SEED PIECE TREATMENT: Damp-off, Seed Rot, Brown Eye Disease, Verticillium Wilt — Use at rate of 10 ozs. per bushel cut seed pieces (16 ozs. or 1 lb. to 100 lbs. of cut seed pieces). Apply in convenient container or by duster attachment over belt. Coat each seed evenly with a light film of dust. Treat pieces within 6 hours after cutting. If planting is delayed, store treated seed pieces in open crates two days before bagging.

DIRECTIONS FOR USE

POTATO SEED-PIECE: Cutting and Treating: Before planting, cutting machines or knives, trays, tables, barrels or sacks, equipment, trucks and planters should be thoroughly cleaned and sanitized. If not sprouted, subject seed potato tubers to temperatures of 50° to 60° F. for about 10 to 15 days before handling them.

CAPTAN 10 DUST

POTATO SEED TREATER



CAUTION

Do not allow this material to drift onto neighboring crops or use in a manner or at a time other than in accordance with directions, because plant injury, excessive residues or other undesirable results may occur.

Do not use treated seed for food or feed purposes.

May be harmful if swallowed or inhaled. May cause irritation of eyes, nose, throat and skin! In case of contact, flush with plenty of water; for eyes get medical attention. Wear rubber gloves and respirator when mixing Seed Treater with potato seed pieces.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Dispose of excess treated seeds or wastes by burial.

Do not reuse container. Destroy when empty.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of **MERCHANTABILITY** or **FITNESS FOR A PARTICULAR PURPOSE**, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

AN AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENT:

Captan (N- [(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide) 10%

INERT INGREDIENTS 90%

Not licensed for use in countries where valid patents exist. 100%

E.P.A. Reg. No. 11682-5
SUB REG. NO. 476-655-15154

CAUTION Keep Out of Reach of Children

HARMFUL IF TAKEN INTERNALLY

AVOID INHALING DUST

AVOID CONTACT WITH SKIN!

AVOID CONTAMINATION OF FEED AND FOOD

STORE IN A COOL DRY PLACE; PROTECT FROM EXCESSIVE HEAT

A PRODUCT OF

SIM-CHEM

FERTILIZERS AND CHEMICALS DIVISION • J. R. SIMPLOT COMPANY

Post Office Box 810

Mountain Home, Idaho 83647

NET WEIGHT 50 LBS.

RECOMMENDATIONS

POTATO (SEED PIECE) TREATMENT: Damp-off, Seed Rot, Brown Eye Disease, Verticillium Wilts — Use at rate of 10 ozs. per bushel cut seed pieces (16 ozs. or 1 lb. to 100 lbs. of cut seed pieces). Apply in convenient container or by duster attachment over belt. Coat each seed evenly with a light film of dust. Treat pieces within 6 hours after cutting. If planting is delayed, store treated seed pieces in open crates two days before bagging.

DIRECTIONS FOR USE

POTATO SEED-PIECE: Cutting and Treating: Before planting, cutting machines or knives, trays, tables, barrels or sacks, equipment, trucks and planters should be thoroughly cleaned and sanitized. If not sprouted, subject seed potatoes to temperatures of 50° to 60° F. for about 10 to 15 days before handling them.