




KELTHANE 4-DUST

AN AGRICULTURAL MITICIDE



OBSERVE FOLLOWING USE PRECAUTIONS

Apply when mites first appear and repeat at 7-10 day intervals, or only as needed to maintain control, but do not use closer to harvest than specified.

Kelthane 4 percent dusts may be applied with ground or air equipment. Make applications when air is quiet, usually early morning or evening. Use the higher recommended rates for air application. Do not make aerial application when ground wind exceeds 4 to 5 miles per hour.

The grower is responsible for residues on his crops as well as for damages caused by drift from his property to others.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

ACTIVE INGREDIENT:

1,1-Bis(p-chlorophenyl)-2,2,2-trichloroethanol 4.0%

INERT INGREDIENTS

96.0%

KELTHANE is a trademark Reg. in the U.S. Pat. Off. by Rohm & Haas Company
E.P.A. Reg. No. 11682-13 100.0%

CAUTION

Keep Out of Reach of Children

HARMFUL IF TAKEN INTERNALLY

AVOID INHALING DUST

AVOID CONTACT WITH SKIN

AVOID CONTAMINATION OF FEED AND FOOD

STORE IN A COOL DRY PLACE

PROTECT FROM EXCESSIVE HEAT

A PRODUCT OF

SIM-CHEM

MINERALS AND CHEMICALS DIVISION • J. R. SIMPLOT COMPANY

Post Office Box 810

Mountain Home, Idaho 83647

NET WEIGHT 50 LBS.

RECOMMENDATIONS

HOPS: To control two-spotted mite — use 4 percent Kelthane Dust at 20 to 30 pounds per acre during early season. Increase dosage to 25 to 35 pounds per acre for mid to late season applications. Do not use within 7 days of harvest.

ALFALFA, CLOVER OR OTHER LEGUMES FOR SEED: To control two-spotted, Pacific, carmine and strawberry (Atlantic) mites — use 25 to 35 pounds of 4 percent Kelthane Dust per acre.

Use restricted to California, Idaho, Oregon, Nevada and Washington. Crop cuttings not to be fed or baled. The fields must be cleared after harvest by burning or by discing into the soil.

BEANS (dry and succulent): To control two-spotted, carmine, Pacific, and strawberry (Atlantic) mites — use 20 to 30 pounds of 4 percent Kelthane Dust per acre. Do not feed straw or crop residues to meat or dairy animals. Do not apply within 7 days of harvest.

APPLES, PEARS, CRABAPPLE, QUINCES: To control European red, two-spotted, McDaniel, clover (brown almond), Schoene, Willamette, apple rust mites — use 40 to 50 pounds of 4 percent Kelthane Dust per acre. Do not apply within 7 days of harvest.

PEACHES, APRICOTS, NECTARINES, CHERRIES, PLUMS, PRUNES, FIGS: For European red, clover (brown almond), two-spotted, Schoene, McDaniel, Pacific, Willamette, peach silver, plum nursery and yellow mites — use 40 to 50 pounds 4 percent Kelthane Dust per acre. To avoid excessive residues do not repeat application within 30 days, and do not use within 14 days of harvest.