




SIM-CHEM

P.S.T. 6 POTATO SEED TREATER

ACCEPTED



CAUTION

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Dispose of excess treated seeds or wastes by burial. Do not reuse container. Destroy when empty.

Do not allow this material to drift onto neighboring crops or use in a manner or at a time other than in accordance with directions, because plant injury, excessive residues or other undesirable results may occur.

May be harmful if swallowed or inhaled. May cause irritation of eyes, nose throat and skin! In case of contact, flush with plenty of water; for eyes get medical attention. Wear rubber gloves and respirator when mixing Seed Treater with potato seed pieces.

POTATO SEED-PIECE:

Dosage and Application: Use 1 lb. of 6% (active ingredient) dust per 100 lbs. of cut seed-pieces. Apply it so that the entire surface of the seed piece is thoroughly covered by the fungicide dust. Place treated seed in clean, disease-free containers or

ACTIVE INGREDIENTS: A coordination product of zinc ion manganese ethylene bisdithiocarbamate 6.00%
In which the ingredients are:
Manganese++ 1.20%
Zinc++ 0.15%
Ethylene bisdithiocarbamate ion (C₂H₂N₂S₂) 4.65%

INERT INGREDIENTS 94.00%
Total 100.00%

E.P.A. Reg. No. 11682-15

EPA Est. No. 08917/D01

**CAUTION Keep Out of Reach of Children
HARMFUL IF TAKEN INTERNALLY**

AVOID INHALING DUST – AVOID CONTACT WITH SKIN

AVOID CONTAMINATION OF FEED AND FOOD

STORE IN A COOL DRY PLACE

PROTECT FROM EXCESSIVE HEAT

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

A PRODUCT OF

SIM-CHEM

MINERALS AND CHEMICALS DIVISION • J. R. SIMPLOT COMPANY

Post Office Box 810

MOUNTAIN HOME, IDAHO 83647

SCHEM 38

NET WEIGHT 50 LBS

DIRECTIONS FOR USE – Continued

store in bulk at proper temperature and humidity. Prevent inhalation of dust by use of exhaust fans during treatment.

Cutting and Treating: Before planting, cutting machines or knives, trays, tables, barrels or sacks, equipment, trucks and planters should be thoroughly cleaned and sanitized. If not sprouted, subject seed potatoes to temperatures of 50 to 60 F. for about 10 to 15 days before handling them.

- A. Controls Fusarium seed-piece decay, the most common seed and soil borne fungus disease of potatoes.
- B. Protects seed which is held for planting.

The cut and treated seed may be planted immediately. If planting is delayed, follow the recommended storage practices of cut seed pieces in your area.

Do not use treated seed for food or feed purposes.