

P.S.T. 6

POTATO SEED TREATER

ACCEPTED

NOV 24 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the use of P.S.T. 6 on potatoes.
EPA Reg. No. 1682-15

ACTIVE INGREDIENTS: A coordination product of zinc ion manganese ethylene bisdithiocarbamate 6.00%

In which the ingredients are:

Manganese+++ 1.20%
Zinc++ 0.15%
Ethylene bisdithiocarbamate ion (C₂H₄N₂S₂) 4.65%

Other INGREDIENTS: 94.00%
TOTAL 100.00%

EPA Reg. No. 11682-15

EPA Est. No. 8917-ID-1

CAUTION

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

HARMFUL IF TAKEN INTERNALLY

AVOID INHALING DUST—AVOID CONTACT WITH SKIN

May be harmful if swallowed or inhaled. May cause irritation of eyes, nose, throat and skin. Do not rub eyes or mouth with hands. Keep all unprotected persons, children, livestock and pets away from treated area or where there is a danger of drift.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF ON SKIN wash with soap and water. If irritation persists, get medical attention.

IF IN EYES flush with water for at least 15 minutes. Get medical attention.

IF INHALED remove victim to fresh air. Apply respiration if indicated.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Cover or incorporate spilled treated seed. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not allow this product to drift onto neighboring crops or use in a manner different from the label directions.

PERSONAL PROTECTIVE EQUIPMENT

For seed piece treatment workers and workers planting seed. HANDLE THIS PRODUCT ONLY WHEN WEARING THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT: a long-sleeve shirt and long pants or a coverall that covers all parts of the body except the head, hands, and feet; chemical resistant gloves; shoes, socks, and goggles or a face shield. Dunning mixing and loading, a chemical resistant apron must also be worn.

IMPORTANT! Before removing gloves, wash them with soap and water. Always wash hands, face, and arms with soap and water before eating, smoking or drinking. Always wash hands and arms with soap and water before using the toilet.

After work take off all clothes and shoes. Shower using soap and water. Wear only clean clothes. Do not use contaminated clothing. Wash protective clothing and protective equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with mancozeb must be destroyed according to state and local regulations.

HEAVILY CONTAMINATED OR DRENCHED CLOTHING CANNOT BE ADEQUATELY DECONTAMINATED.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

POTATO SEED-PIECE TREATMENT

Dosage and Application: Use 1 lb. of 6% (active ingredient) dust per 100 lbs. of cut seed pieces. Apply it so that the entire surface of the seed-piece is thoroughly covered by the fungicide dust. Place treated seed in clean, disease-free containers or store in bulk at proper temperature and humidity. Prevent inhalation of dust by use of exhaust fans during treatment.

Cutting and Treating: Before planting, cutting machines or knives, trays, tables, barrels or sacks, equipment, trucks and planters should be thoroughly cleaned and sanitized. If not sorouted, subject seed potatoes to temperatures of 50 degrees to 60 degrees F. for about 10 to 15 days before handling them.

A. Controls Fusarium seed-piece decay, the most common seed-and-soil-borne fungus disease of potatoes.

B. Protects seed which is held for planting.

The cut and treated seed may be planted immediately. If planting is delayed, follow the recommended storage practices of cut seed pieces in your area.

Do not use treated seed for food or feed purposes.

NOTICE: This product will not control bacterial decay such as blackleg and soft rot, or diseases present within the seed.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

STORAGE: Store in a cool, dry place. Avoid contact with floor and moisture.

DISPOSAL: Completely empty bag into application equipment. Then dispose of in a sanitary landfill, or by incineration, or if allowed by State and local authorities, by burning. If burned stay out of smoke.

DISCLAIMER - J. R. Simplot Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the DIRECTIONS FOR USE. It is impossible to eliminate all risks associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of factors such as weather conditions, presence of other materials, soil conditions, or the manner of use or application, all of which are beyond the control of J. R. Simplot Company. All such risks shall be assumed by the user. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values or any other special or indirect damages. J. R. Simplot Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY.

AGC 594 (8/87)

NET CONTENTS 50 LBS.

Simplot

J. R. Simplot Company
Agricultural Chemical Department
P.O. Box 198, Lathrop, CA 95330
P.O. Box 15057, Boise, ID 83715

BEST AVAILABLE COPY