

SIM-TEC 0.50 POTATO SEED TREATER

ACTIVE INGREDIENT:		By Wt.
'2-(4-thiazoly)-benzimidazole _____		0.50%
INERT INGREDIENTS: _____		99.50%
	TOTAL	100.00%

* Thiabendazole, a product of MSD AGVET,
Div. of Merck & Co., Inc.
Rahway, New Jersey 07065, USA

EPA Reg. No. 11682-30

EPA Est. No. 8917-ID-1

KEEP OUT OF REACH OF CHILDREN

CAUTION


STATEMENT OF PRACTICAL TREATMENT

FIRST AID—If in eyes or on skin: Flush with water; for eyes, get medical attention.

IF SWALLOWED—Call a physician or poison control center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

SEE BACK PANEL

FOR ADDITIONAL PRECAUTIONARY STATEMENTS



PRECAUTIONARY STATEMENT
HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION

Do not get in eyes or on skin. Wear goggles and rubber gloves when handling. Avoid breathing dust. Wear a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provision of 30 CFR Part 11. Wash thoroughly after using. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

GENERAL: For use on Russet Burbank variety only. SIM-TEC 0.50 fungicide dust applied to the cut potato seed pieces protects seed from invasion by FUSARIUM decay fungi present in the soil or on the seed surface and suppresses the development of RHIZOCTONIA STEM CANKER. The use of high quality seed carefully selected and handled, is the first consideration for obtaining full stands.

SIM-TEC 0.50 will not control bacterial decay such as blackleg and soft rot, or diseases present within the seed.

SEED TREATMENT: Russet Burbank variety. For protection against FUSARIUM seed decay and suppression of RHIZOCTONIA STEM CANKER. Use SIM-TEC 0.50 at the rate of 1 lb. per 100 lbs. of cut seed. Apply immediately following cutting with dust dispensing equipment which will give a uniform deposit over the entire seed piece. Treated seed should be planted as soon as possible in warm moist soil. Where storage is necessary, seed should be stored in disease-free holding areas under moderate conditions of humidity and temperature, with adequate ventilation.

Do not use treated seed pieces for food or feed purposes.

20F4
STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

STORAGE: Store in a cool, dry place. Avoid contact with floor and moisture.

DISPOSAL: Completely empty bag into application equipment. Then dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

NET CONTENTS 50 LBS.

Simplot

J. R. Simplot Company
Agricultural Chemical Department
P.O. Box 198, Lathrop, CA 95330
P.O. Box 15057, Boise, ID 83715

POTATO SEED TREATER

ACTIVE INGREDIENT:	By Wt.
*2-(4-thiazoly)-benzimidazole _____	0.50%
INERT INGREDIENTS: _____	99.50%
TOTAL	100.00%

*Thiabendazole, a product of MSD AGVET,
Div. of Merck & Co., Inc.
Rahway, New Jersey 07065, USA

EPA Reg. No. 11682-30

EPA Est. No. 8917-ID-1

KEEP OUT OF REACH OF CHILDREN


CAUTION

STATEMENT OF PRACTICAL TREATMENT

FIRST AID—If in eyes or on skin: Flush with water; for eyes, get medical attention.

IF SWALLOWED—Call a physician or poison control center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

**SEE BACK PANEL
FOR ADDITIONAL PRECAUTIONARY STATEMENTS**



CAUTION

Do not get in eyes or on skin. Wear goggles and rubber gloves when handling. Avoid breathing dust. Wear a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provision of 30 CFR Part 11. Wash thoroughly after using. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

GENERAL: For use on Russet Burbank variety only. SIM-TEC PLUS fungicide dust applied to the cut potato seed pieces protects seed from invasion by FUSARIUM decay fungi present in the soil or on the seed surface and suppresses the development of RHIZOCTONIA STEM CANCKER. The use of high quality seed carefully selected and handled, is the first consideration for obtaining full stands.

SIM-TEC PLUS will not control bacterial decay such as blackleg and soft rot, or diseases present within the seed.

SEED TREATMENT: Russet Burbank variety. For protection against FUSARIUM seed decay and suppression of RHIZOCTONIA STEM CANCKER. Use SIM-TEC PLUS at the rate of 1 lb. per 100 lbs. of cut seed. Apply immediately following cutting with dust dispensing equipment which will give a uniform deposit over the entire seed piece. Treated seed should be planted as soon as possible in warm moist soil. Where storage is necessary, seed should be stored in disease-free holding areas under moderate conditions of humidity and temperature, with adequate ventilation.

Do not use treated seed pieces for food or feed purposes.

Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

STORAGE: Store in a cool, dry place. Avoid contact with floor and moisture.

DISPOSAL: Completely empty bag into application equipment. Then dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

NET CONTENTS 40 LBS.

Simplot

J. R. Simplot Company
Agricultural Chemical Department
P.O. Box 198, Lathrop, CA 95330
P.O. Box 15057, Boise, ID 83715