

11685-19

2/26/2002

1/3



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

FEB 26 2002

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Theodore D. Head
Nufarm, UK
c/o Nufarm Americas Inc.
1333 Burr Ridge Parkway, Suite 125A
Burr Ridge, IL 60521

Dear Mr. Head:


Subject: Add Use on Triticale
Nufarm Rhomene MCPA Amine Herbicide
EPA Registration No. 11685-19
Your Submission Dated February 7, 2002

The amendment referred to above, submitted in connection with registration under section 3(c)(7)(A) of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) or 4(a) when the Agency requires all registrants of similar products to submit such data.
2. Submit one (1) copy of your final printed labeling before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records. If you have any questions concerning this letter, please contact Mr. James Stone at 703-305-7391.

Sincerely yours,


Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL BOOKLET FOR NUFARM RHOMENE MCPA AMINE HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Nufarm Rhomene and this supplement.



Rhomene MCPA **ACCEPTED**
Amine Herbicide **with COMMENTS**
In EPA Letter Dated:

FEB 26 2002

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 11685-19

EPA Reg. No. 11685-19

TRITICALE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

All applicable directions and precautions in the Nufarm Rhomene label booklet must be followed.

Also refer to additional "GENERAL INFORMATION" and "WEEDS CONTROLLED" in the label booklet. INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY, DO NOT USE THIS PRODUCT. Product should not be used in or near greenhouses.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR, Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

RECOMMENDED RATES OF PRODUCT PER ACRE*

Weeds	RATE	Directions
Triticale(not underseeded with legumes)		
Annual Weeds	1/2 pint	Apply after grain is in the 4 leaf stage but not forming joints in the stem. Do not spray in the boot to dough stage.
Annual and biennial broadleaf weeds	1 to 2 pints*	Apply after grain is fully tillered (usually about 4

Perennial broadleaf weeds	1- 1/2 to 2 pints*	to 8 inches high) but not forming joints in the stem. Do not spray grain in boot to dough stage.
Emergency Weed Control in Triticale		Apply when weeds are approaching bud stage, but do not spray grain during the boot to dough stage. The 3 pints (1-1/2lb. acid equivalent) per acre application can produce injury to wheat. Balance the severity of your weed problem against the possibility of crop damage. Where perennial weeds are scattered, spot treatment is suggested to minimize the extent of crop injury.
Perennial broadleaf weeds	3 pints*	
Triticale underseeded with legumes	1/2 pint	Treat after grain is tillered and in the 4-leaf stage and when legume seedlings are 2 to 3 inches tall. Small grains should form a protective canopy over the legume seedlings. Do not use more than 6 gallons of water per acre; higher volumes may result in injury to legumes. Do not apply to grain underseeded with vetch or sweet clover. Do not spray when the grain is in the boot to dough stage.
Alfalfa, alsike, red and ladino clovers, trefoil.		

* Use the lower rate if small annual and biennial weeds are the major problem. Use the higher rate if perennial weeds or annual and biennial weeds are present which are in the hard-to-kill categories as determined by local experience. The higher rates increase the risk of grain injury and should be used only where the weed control program justifies the risk of grain damage. Do not apply to grain in the seedling stage.

NOTICE

Before buying or using this product, read "WARRANTY LIMITATIONS AND DISCLAIMER" in the label booklet. If terms are not acceptable, return unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by the Buyer or any other User constitutes acceptance of the terms under the "WARRANTY LIMITATIONS AND DISCLAIMER".

**Manufactured For:
Nufarm Inc.
St. Joseph, MO 64504**