WHERE TO USE

WEED-RHAP A-6D is used to control broad-leaved weeds in cereal crops, corn, sorghum, flax, weeds and brush in pastures, turf rights of way, similar noncrop uses, and for aquatic weed control.

PLANTS CONTROLLED

WEED-RHAP A-6D will kill or control the following as well as many other noxious plants susceptible to 2.4 D

Puncturevine Arrowhead Purslane Ground ivy Artichoke Bindweed hedge, field. Rush Russian thistle Hoary cress and European) Sagebrush Honeysuckle Bitter wintercress Shepherdspurse Indigo Boxelder Smartweed Ironweed Buckhorn Jimsonweed Sowthistle **Bull thistle** Lambsquarters Stinkweed Bulrush Locoweed Sumac Burdeck Sunflower Mexicanweed Bur ragweed Virginia creeper Morningglory Buttercup Mustard Canada thistle Waterlily Catnip Nutgrass Waterprimrose Parrotfeather Chickweed Wild garlic Pennywort Chicory Cocklebur Wild lettuce Pigweed Wild onton Coffeebean Plantain Wild radish Poison ivy Creeping Jenny Pokeweed Curly indigo Elderberry Witchweed Povertyweed

DIRECTIONS FOR USE


- Mix the recommended amount of product with sufficient water to obtain good spray coverage. Do not mix this product with oil. Best results are obtained when applications are made while the weeds are young and actively growing.
- On Fall-Planted Wheat, Oats, and Barley Use 3/2 to 1 pint of product in 5 to 10 gallons of water to cover one acre. Apply in early spring when weeds are small and before the crop has reached the boot stage.
- On Spring-Planted Wheat, Oats, and Barley—Use 3/3 pint in 5 to 10 gallons of water to cover one acre. Apply after the fully tillered stage, except during the boot to dough stage.
- DO NOT forage or graze treated grain fields within two weeks after treatment with 2,4-D. DO NOT feed treated straw to livestock.
- On Corn and Sorghum—Use 3/3 pint in 5 to 10 gallons of water to cover one acre when weeds are in active growth. Local climatic condition cetermines when treatment should be made. Best results are usually obtained when weeds are small and corn is 4 to 18 inches tall. Do not apply from tasseling to dough stage. Treat sorghum when 4 to 12 inches tall. Use drop nozzles when either crop is over 10 inches tall.
- On Flax—Use 1/3 pint in 10 to 15 gallons of water to cover one acre. Apply when flax is 3 inches high to first bud stage. Do not apply after the flax has formed flower buds.
- On Rice—Use 3/3 to 1 quart in 5 to 10 gallons of water to cover one acre when weeds are in active growth stage. Rice plants are sensitive to 2,4-D in early stages of growth; therefore, it is advisable to delay spraying until the second or third week after flooding. Water in the field should be shallow enough to permit direct application of the spray material to the weeds. Make all treatments well in advance of heading.
- For Aquatic Weeds in lakes, ponds, drainage ditches, and marshes—Use 1% to 3 pints of product in 50 to 100 gallons of water per acre. Apply in enough water to wet foliage thoroughly. Application should be made when leaves are fully developed above water line and plants are actively growing.

 DO NOT apply to more than 1/3 to 1/2 of a lake or pond in any one month, because excessive decaying vegetation may deplete oxygen content of water, killing fish.
- On Pastures, Ornamental Turf, and Noncrop Areas—Use 3/3 to 11/3 quarts and enough water to give sufficient coverage to one acre on established stands of perennial grasses, depending on type of weeds and stage of growth. Do not use on creeping grasses such as bent except for spot spraying. Newly seeded turf should not be treated until after the second mowing and at the lower dosage rate.
- DO NOT graze dairy animals on treated areas within seven days after application.
- Perennial and Other Hard-to-Control Weeds may require a repeat application to give adequate control.



HERCULES

WEED-RHAP®



2,4-D

6 LBS. OF 2,4-D ACID EQUIVALENT PER GALLON

CAUTION: KEEP OUT OF REACH OF CHILDREN.
READ SIDE PANEL FOR ADDITIONAL HANDLING PRECAUTIONS.

ACTIVE INGREDIENT:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid (Equivalent to 57.4% of 2,4-Dichlorophenoxy-

acetic Acid)
INERT INGREDIENTS:

30.9%

TOTAL 100.0%

MADE BY

AGRICULTURAL CHEMICALS
SYNTHETICS DEPARTMENT

HERCULES INCORPORATED

WILMINGTON, DELAWARE, U.S.A.

WARNING

- leaved plants that are desirable such as cotton, soybeans, tomatoes, grapes, clover, ornamentals. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth.
- Minute quantities may cause severe injury by drift. Avoid drift of spray and use a coarse spray which is less likely to drift.
- DO NOT contaminate water in irrigation ditches or water used for domestic purposes.
- Users should note that herbicide treatment of public waters requires a permit from appropriate state agencies in most states. Your State Conservation Department or Game and Fish Commission will aid you in securing a permit if required in your state.
- DO NOT store this product near fertilizers, seeds, insecticides, or fungicides. DO NOT contaminate feed or foodstuffs.
- DO NOT use spray equipment contaminated with this product for any other purpose. DO NOT apply insecticides, fungicides, or other agricultural products unless the equipment has been cleaned very carefully with a suitable chemical cleaner.
- If stored below freezing, it may be necessary to warm to 40°F, and agitate before using. This does not affect the efficiency of the product.
- DO NOT mix with liquid fertilizers.
- DO NOT reuse empty containers. Destroy by burying in an isolated place away from water supplies.

CAUTION

This product may cause skin irritation. Avoid inhaling spray mist. Do not take internally. Avoid contact with the eyes, skin, or clothing. Flush eyes with clear water and get prompt medical attention.

Keep out of reach of children.

H10H1 - 1H111

NONWARRANTY

Hercules warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purpose. Since timing, method of application, weather, plant, and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

THE NOTE OF SEA

WHEFE TO USE

 $(a,b) = (a,A) \cdot (A + b) \cdot (a + b)$

PLANTS CONTROLLED

WELL FOR APPARED A SECTION OF INTERPRETATION OF A SECTION AND ASSESSED.

	**	• • • • • •
	•	. ••
•	* ***	•
••		
		* * A
•		•
* * * * * *		
· · ·		A
	•••	• • •
		10 mm
-	A Company of	:. ·
••, . • .	• - · ·	•
April 19 September 1995	F 8	<u> 444</u>
* . ***** *	Property (ή Α Συνου
f · · · · ·	for the second	AN AND WHEEL .

DIRECTIONS FOR USE


- Mix the recommended amount of product with sufficient water to core good spray average. It got may the product with general some brinds when painted is a compate when the word over a good set of actively gray K.
- On Fall Planted Wheat, Oats, and Barley Use 23 to 1 pint of production to 1 kg in 5 th Author to 2 cm in the 1 miles of the control of the
- On Spring Planted Wheat, Oats, and Barley -- Use Figure 10 to 10 t
- DO NOT forage or graze treated grain fields within two weeks after
- Or Corn and Sorghum Use pint in 5 to 10 gallons of water to
- On Flax Use ... punt in 10 to 15 gallons of water to cover one ... All cover one to some they represent the first of the cover one ...
- Rice Use 2, to 1 quart in 5 to 10 gallons of water to cover the active wholes one mark to go with state. Remove to the active with state in Remove to the active with state in Remove to the active with the active with the active for the active for the active for the active for the active with the act
- For Aquatic Weeks, is lakes, ponds, drainage ditches, and marshes to the state of t
- Pastures Organierial Furf and Noorrop Areas. Use or to 1
- the MCC containing a second of the Both containing with the containing

Control of the Contro



HERCULES

WEED-RHAP®



CAUTION: KEEP OUT OF REACH OF CHILDREN.
READ SIDE PANEL FOR ADDITIONAL HANDLING PRECAUTIONS.

ACTIVE INGREDIENT:
Limethylamine Salt of 2.4:
Dichlorophenoxyacetic Acid 69 1%

(Equivalent to 57.4% of 2,4 Dichlorophenory acetic Acid)

INTRI HIGHEDIENTS 30 9° 0

TOTAL 100.0%

MYDEBY

AGRICULTURAL CHEMICALS
SYNTHETICS DEPARTMENT

HERCULES INCORPORATED

WILMINGTON DELAWARE U.S.A.

WARNING

- DO NOT apply this product to susceptible broad with the control of the control of
- Minute quantities may cause severe injury by define A. The first stray and use the stray and use the stray and use the second of the second of
- DO NOT contaminate water in irrigation ditches to a the contaminate water in irrigation ditches
- Users should note that herbicide treatment of public waters in a test of permit from applicate state of the estimate of states. Your State Conservation Department or Game and fish Conservation will aid you in security or remort frequency in your state.
- DO NOT store this product near fertilizers, seeds, insecticides, or fungicides. DO NOT contaminate feed or foodstuffs.
- DO NOT use spray equipment contaminated with this product for any other purpose. DO N apply inserticides funciodes, or other aim cultural products unless the equipment has been cleaned very carefully with a suitable of emical cleaner.
- If stored below freezing, it may be necessary to warm to 4% found apitate before using 11% does not affect the efficiency of the product.
- DO NOT mix with liquid fertilizers.
- DO NOT reuse empty containers. Destroy by buryme in an incident place and, from water sin lines.

CAUTION

- This product may cause skin irritation. Avoid inhaling spray must. Do not take internally. Accordant contact with the eyes, skin or fothers. Fust eyes, with clear water and yet prompt medical attention.
- Reep out of reach of children

NONWARRANTY

Hercules warrants that this herbicide conforms to the chem at description on its tabel. When used a exceedance with taber directions unless mero a conditions this herbicide is responding to this intended purpose. Since the normal method of applie at an weather plant or to condition menture with other chemicals. The conditions affecting the use of this product on the rich error te, and our control as according to the discontrol, the area to be to a conditions which are according to the product control, the according of the product control, the according of the product control and according to the according to the product control of the