


(FRONT PANEL)



TRANSVAAL  
TRANSMINER<sup>R</sup> OA-3D  
HERBICIDE  
2,4-D

3 Lbs. of 2,4-D Acid Equivalent Per Gallon

ACTIVE INGREDIENTS:

N,N-Dimethyl Oleyl-Linoleyl Amine Salt of 2,4-Dichloro- phenoxyacetic Acid (Equivalent to 35.2% or 3 lbs. 2,4-D per gallon)	61.5%
---	-------

INERT INGREDIENTS:

	38.5%
TOTAL	100.0%

EPA REGISTRATION NO. 11687-41

MADE BY

TRANSVAAL, INC.

JACKSONVILLE, ARKANSAS 72076

CAUTION: KEEP OUT OF REACH OF CHILDREN  
READ SIDE PANELS FOR ADDITIONAL HANDLING PRECAUTIONS.

NET CONTENTS: GAL.

WHERE TO USE

TRANSMINER<sup>R</sup> OA-3D controls broadleafed weeds in cereal crops, corn, sorghum, sugar cane; weeds and brush in rangeland, pastures, rights-of-way, turf and similar non-crop uses. For use in pear and apple orchards.

PLANTS CONTROLLED

TRANSMINER<sup>R</sup> OA-3D will kill or control the following as well as many other noxious plants susceptible to 2,4-D:

Arrowhead	Chicory	Lamb's-quarters	Rush
Artichoke	Cocklebur	Locoweed	Russian thistle
Bindweed (hedge, field and European)	Coffeebean	Mexicanweed	Sagebrush
Bitter wintercress	Creeping jenny	Morning-glory	Shepherd's-purse
Boxelder	Curly indigo	Mustard	Smartweed
Buckhorn	Duckweed	Nutgrass	Sowthistle
Bull thistle	Elderberry	Parrotfeather	Stinkweed
Bulrush	Goldenrod	Pennywort	Sumac
Burdock	Ground ivy	Pigweed	Sunflower
Bur ragweed	Hemp	Plantain	Virginia creeper
Buttercup	Hoary cress	Poison ivy	Wild garlic
Canada thistle	Honeysuckle	Pokeweed	Wild lettuce
Catnip	Indigo	Povertyweed	Wild onion
Chickweed	Ironweed	Puncturevine	Wild raddish
	Jimsonweed	Purslane	Willow

DIRECTIONS FOR USE

Add the recommended amount of product to an approximately equal volume of water and STIR THOROUGHLY. Add this mix to the remaining volume of water to be used in the application. Agitate until spray mixture is uniform.

Best results are obtained when applications are made while the plants are well-leaved and actively growing.

On Fall-Planted Wheat, Oats and Barley - Use 2/3 to 1 pint of product in 5 to 10 gallons of water to cover one acre. Apply in early spring when weeds are small and before the crop has reached the boot stage.

On Spring-Planted Wheat, Oats and Barley - Use 2/3 pint in 5 to 10 gallons of water to cover one acre. Apply after the fully tillered stage, except during the boot to dough stage.

DO NOT forage or graze treated grain fields within 2 weeks after treatment with 2,4-D. DO NOT feed treated straw to livestock.

On Corn and Sorghum - Use 1 1/2 pint in 5 to 10 gallons of water to cover one acre when weeds are in active growth. Local climatic condition determines when treatment should be made. Best results are usually obtained when weeds are small and corn is 4 to 18 inches tall. DO NOT apply from tasseling to dough stage. Treat sorghum when 4 to 12 inches tall. Use drop nozzles when either crop is over 10 inches tall.

For Pre-emergence on Corn - Use 2 1/2 to 5 pints in 5 to 10 gallons of water per acre. Use the higher rate on heavy soils. DO NOT use on sandy soils. Plant corn as deep as feasible.

On Rice - Ground Application - Dilute 1 to 1 1/2 quarts of product in 5 to 10 gallons of water and apply this volume to one acre.

Aerial Application - Dilute 1 to 1 1/2 quarts of product in 4 to 7 gallons of water and apply this volume to one acre.

Applications should be made between tillering and boot stages. Fields should be drained during application.

On Sugar Cane - Use 1 1/3 to 2 2/3 quarts of product in 5 to 25 gallons of water per acre as a pre- and/or post-emergence spray. Consult local agricultural experiment station or extension service weed specialist on specific use of this product alone or in combination with other chemicals.

For Lawn, Golf Courses and Fairway Surf - Apply 2 to 4 pints per acre in sufficient water to provide adequate coverage in spring or fall when weeds are actively growing. DO NOT apply to newly seeded turf until sod is formed and grass has been cut several times. Injury may result if applied to grasses such as St. Augustine, Bent and Dickcandra. Clover may also be injured. Bare areas can be reseeded in 3 to 6 weeks.

On Rangeland, Pastures, and for General Weed Control (rights-of-way, fence rows and other noncrop areas) - Use 1 1/4 to 5 pints of product in sufficient water to give good coverage to one acre, depending on type of weeds and state of growth.

Use only on established stands of perennial grasses.

DO NOT graze dairy animals on treated pastures within 7 days after application.

For Use in Pear and Apple Orchards - Apply 3 to 5 1/2 pints of product per acre in sufficient water to provide adequate coverage and wetting of weeds. Use higher dosage for perennial weeds. Avoid applications to bare ground. Direct spray at undesirable plants. DO NOT allow spray to contact foliage, stems, fruit or trunks of fruit trees. To minimize drift, spray on still days with low pressure and coarse nozzle.

Perennial and other hard-to-control weeds may require a repeat application to give adequate control.

#### CAUTION

This product may cause skin irritation. Avoid inhaling spray mist. Harmful if swallowed. Avoid contact with the eyes, skin or clothing. In case of contact with eyes, flush with clear water and get prompt medical attention.

DO NOT apply this product to susceptible broadleaved plants that are desirable such as cotton, soybeans, tomatoes, grapes, clover or ornamentals. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth.

Minute quantities may cause severe injury by drift. Avoid drift of spray and use a coarse spray which is less likely to drift.

This herbicide is toxic to fish. Keep out of lakes, streams or ponds. DO NOT apply where runoff is likely to occur. DO NOT apply when weather conditions favor drift from treated areas.

DO NOT contaminate any body of water by direct application, cleaning of equipment or disposal of wastes and containers. Apply this product as specified on this label.

DO NOT store this product near fertilizers, seeds, insecticides or fungicides.

DO NOT contaminate feed or foodstuffs.

DO NOT use spray equipment contaminated with this product for any other purpose.

DO NOT apply insecticides, fungicides or other agricultural products unless the equipment has been cleaned very carefully with a suitable chemical cleaner.

DO NOT mix with liquid fertilizers.

DO NOT reuse empty containers. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

If stored below freezing, it may be necessary to warm to 40°F and agitate before using. This does not affect the efficiency of the product.

#### WARRANTY

Transvaal warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purpose. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.