

TRANSVAAL

WEED-RHAP[®] LV OXY-4 D HERBICIDE

WHERE TO USE

WEED-RHAP LV OXY-4 D is used to control broad-leaved weeds in cereal crops, corn, sorghum, weeds and brush in rangeland, pastures, rights-of-way, similar noncrop uses; and for aquatic weed control.

PLANTS CONTROLLED

WEED-RHAP LV OXY-4 D will kill or control the following as well as many other noxious plants susceptible to 2,4-D.

Almondwood	Chicory	Lecowood	Sagebrush
Arenaceous	Cocklebur	Mesquited	Sapindaceous
Artemisia	Coffeeweed	Morinda-silky	Smartweed
Bittersweet (hedge, fence, and European)	Croton, Jersey	Mustard	Stinkweed
Bitter-mustard	Curly Indigo	Nutgrass	Sumac
Buckwheat	Duckweed	Parennial	SUNFLOWER
Buckwheat	Fiddleberry	Pennycress	VIRGINIA CLEAVER
Buckwheat	Goldenrod	Pigweed	Water hyacinth
Buckwheat	Ground Ivy	Poa-grass	Water lily
Buckwheat	Hemp	Poison ivy	Wild radish
Buckwheat	Hairy grass	Poison oak	Wild lettuce
Buckwheat	Honey mesquite	Poison sumac	Wild onion
Buckwheat	Horsetail	Puncturevine	Wild radish
Buckwheat	Ivy	Purslane	Willow
Buckwheat	Lambquarters	Rush	
Buckwheat	Russian-thistle		

DIRECTIONS FOR USE

Add the recommended amount of product to about one-half the volume of water to be used in spraying. Mix well, then add remaining water and mix until spray mixture is uniform. Agitate thoroughly while spraying to assure uniform emulsion. Best results are obtained when applications are made while the plants are well-leaved and actively growing.

On Fall-Planted Wheat, Oats and Barley—Use ½ to ¾ pint of product in 5 to 10 gallons of water to cover one acre. Apply in early spring when weeds are small and before crop has reached the boot stage.

On Spring-Planted Wheat, Oats and Barley—Use ½ pint of product in 5 to 10 gallons of water to cover one acre. Apply after the fully tilled stage, except during the boot to dough stage.

DO NOT forage or graze treated grain fields within two weeks after treatment with 2,4-D. DO NOT feed treated straw to livestock.

On Corn and Sorghum—Use ½ pint of product in 5 to 10 gallons of water to cover one acre when weeds are in active growth. Local climatic conditions determine when treatment should be made. Best results are usually obtained when weeds are small and corn is 4 to 18½ inches tall. DO NOT apply from tasseling to dough stage. Treat sorghum when 4 to 12 inches tall. Use drop nozzles when either crop is over 10 inches tall.

For Preemergence on Corn—Use 1 to 2 quarts of product in 5 to 10 gallons of water per acre. Use the higher rate on heavy soils. DO NOT use on sandy soils. Plant corn as deep as feasible.

On Ornamental Turf—Depending on type of weeds and stage of growth, use 1 to 3 pints of product and enough water to give sufficient coverage to one acre on established stands of perennial grasses. DO NOT use on creeping grasses such as bent except for spot spraying. Newly seeded turf should not be treated until after the second mowing, and the lower dosage rate should be used. Possible injury may occur to Dichondra and clover.

On Rangeland, Pastures, and General Weed Control—Use 1 to 4 pints of product in sufficient water to give good coverage to one acre, depending on type of weeds and stage of growth. Use only on established stands of perennial grasses. DO NOT graze dairy animals on treated areas within seven days after application.

For Aquatic Weeds in Lakes, Ponds and Drainage Ditchbanks—Use 2½ to 4½ pints of product in 50 to 100 gallons of water and apply to one acre. Application should be made when leaves are fully developed above water line and plants are actively growing. Treatment of dense weed areas can result in oxygen loss from decomposition of dead weeds. This loss can cause fish suffocation. Therefore, to minimize this hazard treat ½ to ⅓ of the water area in a single operation and wait at least 10-14 days between treatments. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated areas.

Perennial and other Hard-To-Control Weeds may require a repeat application to give adequate control.

CAUTION

This product may cause skin irritation. Avoid inhaling spray mist. Harmful if swallowed. Avoid contact with the eyes, skin or clothing. Flush eyes with clear water and get prompt medical attention.

DO NOT apply this product to susceptible broad-leaved plants that are desirable such as cotton, soybeans, tomatoes, grapes, clover or ornamentals. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth.

Minute quantities may cause severe injury by drift. Avoid drift of spray and use a coarse spray, which is less likely to drift. Although this product contains an ester of low volatility, use it with caution where the spreading of vapors can damage nearby susceptible plants.

DO NOT contaminate water in irrigation ditches or water used for domestic purposes.

DO NOT contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Users should note that herbicide treatment of public waters requires a permit from appropriate state agencies in most states. Your State Conservation Department or Game and Fish Commission will aid you in securing a permit if required in your state.

DO NOT store this product near fertilizers, seeds, insecticides or fungicides.

DO NOT contaminate feed or foodstuffs.

DO NOT use spray equipment contaminated with this product for any other purpose.

DO NOT apply insecticides, fungicides or other agricultural products unless the equipment has been cleaned very carefully with a suitable chemical cleaner.

DO NOT reuse containers. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

If stored below freezing, it may be necessary to warm to 40°F. and agitate before using. This does not affect the efficiency of the product.

DO NOT USE, pour, spill or store near heat or open flame.

WARRANTY

TRANSVAAL warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purpose. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

ACCEPTED

11687-58
10-31-73

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

2,4-D Low Volatile
4 LBS. OF 2,4-D ACID EQUIVALENT PER GALLON

ACTIVE INGREDIENTS:

Butoxyethyl Ester of 2,4-Dichlorophenoxyacetic Acid* 65.7%

Inert Ingredients 34.3%

TOTAL 100.0%

*Equivalent to 45.2% or 4 pounds per gallon of 2,4-Dichlorophenoxyacetic Acid.

EPA REGISTRATION NO. 11687-58

MADE BY

TRANSVAAL, INC.

JACKSONVILLE, ARKANSAS 72076

NET CONTENTS:

GAL.

TP

CAUTION: KEEP OUT OF REACH OF CHILDREN.

SEE SIDE PANELS FOR ADDITIONAL HANDLING PRECAUTIONS