TRANSVAL WEED-RHAP®

WHERE TO USE

WEED-RHAP LV-4-MCPA is recommended to control certain weeds in wheat outs barley tye, flax grassland, fence rows and roadsides and other similar non-crop areas.

PLANTS CONTROLLED

WEED-RHAP! LV-4-MCPA is recommended to kill and to control the following, as well as many other noxious plants susceptible to MCPA

Annual Sow Thistle
Beggarticks
Buttercups
Canada Thistle
Chervil
Cocklebur
Dandelions
Dragon Head mint
Field-Pepper Grass
Goats Beard
Hempnette

Kochia
Lambsquarter
Marsh Elder
Miner's lettuce
Mustards
Perennial Sow Thistle
Plantains
Poison Hemlock
Puncturevine
Purslane

Russian Pigweed Shepherdspurse Stinging Nettle Stinkweed Sunflower Vetch White top Wild radish Yellow rocket

DIRECTIONS FOR USE

Ragweed

Mix the recommended amount of product with sufficient water to obtain good spray coverage. Best results are obtained when spraying is done while the weeds are young and actively growing

WHEAT, OATS, BARLEY and RYE (Aerial and surface applications): Use 1/2 to 1 pint per acre for the more susceptible weeds after crop has reached the 3 to 4 leaf stage up to boot stage. Apply as a water mix spray. Use up to 3 pints per acre for the less susceptible weeds after crop has tillered and up to early boot stage. Do not spray from boot stage to dough stage.


FLAX (Aerial and surface applications): Use 1/4 to 1/2 pint per acre. Use sufficient water to give uniform and adequate coverage. Apply only when weeds are up and when flax is 4 to 8 inches high and before it comes into bud. Treatment after early bud stage may result in severe damage. If Canada Thistle is present it may be necessary to go as high as 3/4 pint per acre to prevent seed head product. Some injury to the flax may result

GRASSLAND (Established grassland) (Aerial and surface applications): Use 1 to 3 pints per acre in sufficient water. (10 to 100) gallons, for thorough coverage. Use higher rate for White Top. Canada Thistle, and other less susceptible weeds. Spray perennials in early-bud to full bloom stage-and regrowth in fall, other weeds in spring or fall, when actively growing.

GRASSES (Grown for seed) (Aerial and surface applications): Use 1 to 2 pints per acre, the higher rate where weed stands are heavy. In established grasses apply in spring before head comes into boot and on seedling grass after grass has tillered. NOTE. In weed control in grasses, repeat treatment may be needed for less susceptible weeds. In some areas bent, builfalo, carpet and St. Augustine grasses may also be injured by treatment.

NON-CROP AREAS (Fence rows and roadsides) (Aerial and surface applications): On Canada Thistle. White Top and Meadow Buttercups, for spot treatment, use 1/4 pint to 3 to 4 gallons of water, or 3/4 gallon per acre in sufficient water to give coverage for more extensive areas. Spray to wet weeds thoroughly when in bud to early bloom and again on fall regrowth.

LV - 4 - MCPA HERBICIDE



M.C.P.A. LOW VOLATILE

4 LBS. OF M.C.P.A. ACID EQUIVALENT PER GALLON

ACTIVE INGREDIENT:

INERT INGREDIENTS:

Isooctyl Ester of 2-Methyl-4-Chlorophenoxyacetic Acid*

73.3% 26.7%

TOTAL 100.0%

'(Equivalent to 47.0% or 4 lbs. per gallon of 2-Methyl-4-Chlorophenoxyacetic Acid) MADE BY

TRANSVAAL, INC.

JACKSONVILLE, ARKANSAS 72076

CAUTION

This product is harmful if swallowed. Avoid contact with skin, eyes or clothing. In case of eye contact, flush eyes with plenty of water and get medical attention. Wash exposed skin with plenty of soap and water. Remove and wash contaminated clothing and shoes before re-use. Avoid inhaling spray mist.

DO NOT apply this product to susceptible plants such as vegetables, flowers, grupes, tomatoes, cotton, beans and other legumes, etc.

Minute quantities may cause severe injury by drift of spray, and use a coarse spray which is less likely to drift. Although this product contains material of low volatility, use it with caution where the spreading of vapors can damage nearby susceptible plants.

DO NOT contaminate water in irrigation ditches or water used for domestic purposes. This product is toxic to fish. Keep out of lakes, streams or ponds DO NOT contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DO NOT store or transport this product near fertilizers, seeds, insecticides or fungicides.

DO NOT contaminate feed or foodstuffs

DO NOT use around home or on recreation sites, or similar areas

DO NOT use spray equipment contaminated with this product for any other purpose unless the equipment has been cleaned very carefully with a suitable chemical cleaner.

DO NOT store in freezing temperatures. If allowed to freeze remix before using

DO NOT use or store near heat or open flame.

DO NOT reuse empty container. Destroy by burying in a safe place away from water supplies.

WARRANTY

TRANSVAAL, INC. warrants that this product conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purpose. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use

NET CONTENTS:

GALLONS

EPA REGISTRATION NO. 11687-79 EPA ESTBLISHMENT NO. 11687-AR-

CAUTION: KEEP OUT OF REACH OF CHILDREN.
READ SIDE PANELS FOR ADDITIONAL HANDLING PRECAUTIONS.