

SUPERNEEM™ 4.5-B BOTANICAL INSECTICIDE

An Insecticide for Use on Vegetables, Ornamentals, Trees, Shrubs, Lawns, and Plants, and In and Around Commercial Nurseries, Greenhouses, Interiorscapes, and the Home

Kills/repels a variety of insect pests including whiteflies, Japanese beetles, caterpillars, aphids, gypsy moths, white grubs, chinch bugs, and diamondback moths.

ACTIVE INGREDIENT:

Azadirachtin	4.5%
INERT INGREDIENTS	95.5%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION


Manufactured By: W. R. Grace & Co.-Conn.
7379 Route 32
Columbia, MD 21044

EPA Reg. No. 11688-12

Net Contents: _____

EPA Est. No. _____

See Back Panel for Additional Precautionary Statements



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Avoid contamination of feed and foodstuffs. Avoid breathing spray mist. In case of eye contact, flush eyes with plenty of water. If on skin, wash with soap and water. Wash contaminated clothes before reuse. If irritation persists, get medical attention.

Environmental Hazards:

This product may be hazardous to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of water.

Physical and Chemical Hazards:

Flammable. Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

Storage: Keep in original container. Store in a cool, dry place, away from direct sunlight, feed or foodstuffs. Keep container tightly sealed when not in use. Do not store below 50°F (10°C) or above 95°F (35°C).

Disposal: Rinse water and unused diluted pesticide may be disposed of on-site or in an approved waste disposal facility. Do not reuse empty container. Triple rinse empty containers and offer for recycling or puncture and dispose of in a sanitary landfill or by incineration, if allowed by state and local authorities. If burned, stay out of smoke.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

General

- Botanical Insecticide Concentrate
- Broad Spectrum Insecticide
-) • Broad Spectrum Organic Insecticide
-) • Organic Insecticide
- Multi-Purpose Insecticide
- Multi-Purpose Organic Insecticide
- Kills and/or repels insects/Japanese beetles.
-) • Kills larval stages of insects.
- Organically kills insects.
-) • Organically repels Japanese beetles.
-) • For Indoor and Outdoor Use
- Formulated for interiorscape use.
- Not for use in food-handling areas.
- Shake well before using.
- SuperNeem™ 4.5-B is most effective when applied 2-3 times in succession.
- Spraying directly onto the pest and a longer duration of leaf wetting increase effectiveness. Apply in early to mid-morning or late afternoon. This is particularly important with whitefly adults which are sedentary on the undersides of leaves at these times.

- SuperNeem™ 4.5-B in diluted solution should be maintained at pH between 3-8. Use outside of this pH range may cause product degradation. Spray solutions should be used within several hours of preparation for maximum effectiveness. Do not store diluted solution for later use.
- Do not add adjuvants (spreaders, surfactants, etc.) to SuperNeem™ 4.5-B as it already contains its own surfactant.
- Do not apply to wilted or otherwise stressed plants, or to newly transplanted material prior to root establishment. Do not apply to known spray-sensitive plants without testing.
- SuperNeem™ 4.5-B has been found to be compatible when used in conjunction with most beneficial insects. It is recommended that a small trial be conducted to assure compatibility before using on a large scale.

NONCROP APPLICATIONS

Directions for Interiorscapes, Plants, Landscapes, Ornamentals, Trees, Lawns, and Shrubs

Commercial Use

- For use to control whiteflies, thrips, mealybugs, leafminers, loopers, caterpillars, beet armyworms, and aphids on bedding plants, potted plants, foliage plants, ornamentals, trees, and shrubs in and around greenhouses, commercial nurseries, interiorscapes, and homes.
- For use to control gypsy moths, weevils, psyllids, webworms, hornworms, spruce budworms, and pine sawflies on trees and shrubs on residential and commercial landscapes.
- SuperNeem™ 4.5-B may be used on all listed fruits, vegetables, vegetable transplants, and herbs both inside and outside of the greenhouse.
- For high volume application, dilute SuperNeem™ 4.5-B to a concentration of 2.25 to 4.5 ounces per 100 gallons of water (1/8 to 1/4 teaspoon of SuperNeem™ 4.5-B per gallon of water). Mix thoroughly. Apply 25-40 psi with hand sprayer or 100-200 psi with power sprayer as a fine spray to both leaf surfaces to runoff. Excessive application is unnecessary and should be avoided.
- For low volume application, apply 0.5 pint of SuperNeem™ 4.5-B (equivalent to 8 grams of active ingredient) per acre in sufficient water to provide adequate coverage.

- Sprays may be applied on a preventative 7-day schedule or at the first sign of insect presence. This schedule is effective under low insect pressure. Under high insect pressure, apply every 3-4 days.
- For drench application in greenhouse plantings, use 2.25 ounces per 100 gallons and apply at the rate of 1 quart of diluted solution per square foot of growing media surface. Repeat at 14-day intervals during the growing season.

Tank Mixing

SuperNeem™ 4.5-B Botanical Insecticide Concentrate has been found to be compatible with most commonly used fungicides, insecticides, and fertilizers. Physical compatibility should first be checked by using the correct proportion of products in a small jar test. Growers should then test tank-mix combinations for phytotoxicity on a sample of plants prior to use. This is also recommended with combinations used before as environmental conditions can alter the interaction between compounds. *Due to the wide variation in climatic conditions, cultural practices, and other factors, the user assumes full responsibility for any crop damage or other liability resulting from the use of SuperNeem™ 4.5-B Botanical Insecticide Concentrate in a tank mix combination.* Do not mix SuperNeem™ 4.5-B with oxidizing agents such as bleach, or strong acids and bases as they will destabilize the product.

Home Use

- For use on bedding plants, potted and foliage plants, ornamentals, trees, shrubs, vegetable gardens, and fruit trees in and around homes to control the immature or larval forms of insect pests.
- For use on trees, lawns, and shrubs on residential landscapes to control the immature or larval stages of pests.
- For use as a repellent against ants, cockroaches, and Japanese beetles.
- Apply 0.5-1.0 pint of SuperNeem™ 4.5-B per acre (or 0.17-0.33 ounces per 1000 square feet) using enough spray volume to obtain thorough coverage, usually 50-100 gallons of diluted material per acre (or 1-2 gallons per 1000 square feet). Apply with a hand sprayer as a fine spray to leaf surfaces to run-off. Be sure to coat undersides of leaves as this is where many insects hide. Excessive application is unnecessary and should be avoided. Do not apply to known sensitive plants without a prior small-scale test application.

- Known sensitive plants
 - African Violet flowers
 - Hibiscus flowers
 - Impatiens flowers

Directions for Repelling Japanese Beetles from Rose Plants

- For best results, apply to roses at the first sign of Japanese beetle emergence in early summer at the rate of 0.5 pint of SuperNeem™ 4.5-B per 100 gallons of water.
- SuperNeem™ 4.5-B is more effective when used as a preventative.
- Spray to run-off, making sure to completely cover all parts of the plant, including buds and flowers
- Repeat application weekly or after rainfall.
- More frequent applications may be necessary during periods of rapid plant growth, as new growth that occurs after application is not fully protected.
- Continue applications as long as adult beetles are present.
- Do not spray water directly onto foliage or otherwise wash off the leaves after treatment. This will reduce the effectiveness of the application.
- After initial application, some beetles may be present on foliage but they will not feed on it.

Directions for Lawns and Turf

SURFACE-FEEDING INSECTS

- For use to control cutworms, armyworms, sod webworms, crickets, chinch bugs, leafhoppers, and grasshoppers.
 - Apply 0.5-1.0 pint of SuperNeem™ 4.5-B per acre (or 0.17-0.33 ounces per 1000 square feet) using enough spray volume to obtain thorough coverage, usually 50-100 gallons of diluted material per acre (or 1-2 gallons per 1000 square feet).
 - Apply at first sign of pest emergence or damage. Reapply as necessary. Be sure to treat under shrubs and plants bordering house. Do not apply before it rains. Do not water turf for 2 days after application.

SUBSURFACE-FEEDING INSECTS

- For use to control white grubs (Japanese beetles, European chafers, dung beetles, green June beetles, May/June beetles, annual white grubs, grub beetle, southern masked chafers, etc.) and crane flies.
 - Apply 0.5-1.0 pint of SuperNeem™ 4.5-B per acre (or 0.17-0.33 ounces per 1000 square feet) using enough spray volume to obtain thorough coverage, usually 50-100 gallons of diluted material per acre (or 1-2 gallons per 1000 square feet).
 - Application should be made soon after adults emerge in summer (1-3 weeks after first sign of adults). Turf should be mowed before application. Irrigate turf prior to application. Do not water turf within 24 hours after application. Do not mow turf within 3 days after application.
- For use to control mole crickets.
 - Apply 0.5-1.0 pint of SuperNeem™ 4.5-B per acre (or 0.17-0.33 ounces per 1000 square feet) using enough spray volume to obtain thorough coverage, usually 50-100 gallons of diluted material per acre (or 1-2 gallons per 1000 square feet).
 - For best results, apply when nymphs are small, in the early spring. If necessary reapply at 1-2 week intervals.
- For use to control billbugs.
 - Apply 0.5-1.0 pint of SuperNeem™ 4.5-B per acre (or 0.17-0.33 ounces per 1000 square feet) using enough spray volume to obtain thorough coverage, usually 50-100 gallons of diluted material per acre (or 1-2 gallons per 1000 square feet).
 - Apply in mid to late spring or at first sign of pest emergence or damage. Do not apply before it rains. Do not water turf for 2 days after application. Reapply as necessary. Repeat treatment in early to mid fall to control possible second generation.

DIRECTIONS FOR FOOD CROP APPLICATION

General Directions

- Do not apply this product through any type of irrigation system.
- Use care when applying near streams, ponds, lakes, or other bodies of water. SuperNeem™ 4.5-B should not be applied when weather conditions favor drift or when the likelihood of runoff is high.

Specific Crop Directions

Application Rate:

Apply 0.25-1.0 pint of SuperNeem™ 4.5-B per acre in sufficient water to assure adequate coverage. The low rate should be used as a preventative when pest pressure is low, or if used in conjunction with adulticide products. Otherwise, the high rate should be used.

CITRUS, POME AND STONE FRUITS	
Crops	Pests
Grapefruits	Greenhouse whiteflies
Kumquats	Sweet Potato Whiteflies
Lemons	Mealybugs
Limes	Leafminers
Oranges	Caterpillars
Apples	Aphids
Crabapples	Fruit flies
Pears	Psyllids
Quinces	Thrips
Jujubes	Fall webworms
Apricots	Weevils
Cherries	Gypsy Moths
Nectarines	Leafhoppers
Peaches	Grasshoppers
Plums	
Prunes	

CUCURBITS	
Crops	Pests
Balsam pears	Greenhouse whiteflies
Chinese waxgourds	Sweet potato whiteflies
Cucumbers	Budworms
Gherkins	Fall armyworms
Gourds	Melonworms
Cantaloupes	Pickleworms
Honeydew melons	Rindworms
Mangoes	Squash vine borers
Pumpkins	Leafminers
Squashes	Squash bugs
Watermelons	

BULB AND COLE VEGETABLES	
Crops	Pests
Broccoli	Greenhouse whiteflies
Brussels sprouts	Sweet potato whiteflies
Bok Choy	Cabbage Loopers
Cabbage	Imported Cabbage worm
Cauliflower	Diamondback moth
Chinese Spinach	Aphids
Celery	Cutworms
Collards	Thrips
Cress	Armyworms
Endive	Leafminers
Fennel	Onion maggots
Kale	Hornworms
Kohlrabi	Webworms
Lettuce	Caterpillars
Mustard greens	Weevils
Parsley	
Rhubarb	
Spinach	
Swiss Chard	
Turnip tops	
Garlic	
Leek	
Onions	
Shallots	

LEGUME AND FRUITING VEGETABLES	
Crops	Pests
Beans	Loopers
Chick peas	Colorado potato beetles
Lentils	Leafminers
Peas	Cabbage loopers
Peanuts	Caterpillars
Soybeans	Greenhouse whiteflies
Eggplants	Sweet potato whiteflies
Ground cherries	Armyworms
Peppers	Cutworms
Tomatoes	Soybean loopers
	Hornworms
	Psyllids
	Tomato fruitworms
	Weevils
	Tomato pinworms
	Pepper weevils
	Garden webworms
	Mexican bean beetles
	Bean leaf beetles

ROOT AND TUBER VEGETABLES	
Crops	Pests
Artichokes	Artichoke plume moths
Beets	Colorado potato beetles
Carrots	Leafhoppers
Cassava	Aster leafhoppers
Ginger	Potato flea beetle
Horseradish	Aphids
Parsnips	Greenhouse whiteflies
Potatoes	Sweet potato whiteflies
Radishes	Psyllids
Rutabaga	Cabbage Loopers
Sweet potatoes	Diamondback moths
Tumeric	Hornworms
Turnips	Webworms
Yams	Caterpillars
Yam beans	Armyworms

SMALL FRUITS AND BERRIES	
Crops	Pests
Blackberry	Gypsy moth
Blueberry	Blueberry leafroller
Boysenberry	Loopers
Cranberry	Fruit tree leafroller
Current	Grape berry moth
Dewberry	Oblique banded leafroller
Elderberry	Hornworm
Gooseberry	Armyworm
Grape	Tobacco budworm
Huckleberry	Cherry fruitworm
Loganberry	Green fruitworm
Raspberry	Grape leaf folder
Strawberry	Grapeleaf skeletonizer
Youngberry	Omnivorous leafroller
	Caterpillars
	Mites
	Thrips