DIRECTIONS FOR USE:

ANNUAL WEEDS:

Apply one gallon of Bochlor 17 (diluted with up to 10 gallons of water) per 150 to 200 square feet when plants are young. Thoroughly wet with a sprayer or sprinkling can all parts of the plant, particularly the growing tips.

PERENNIAL WEEDS:

For the best results treat plants during their dormant stage or when nearing maturity. Good results are obtained with Spring application, but Summer applications are not recommended except as a knockdown treatment. Apply by spraying 1 to 2 gallons of Bochlor 17 per 100 square feet. For deep rooted weeds and grass, such as Johnson, Bindweed, Bermuda and Quack grass the heavier rate is recommended. Re-treatment may be necessary to secure adequate control of certain hard-to-kill weeds.

BOCHLOR 17

NON-INFLAMMABLE NON

DESTROYS:

Quack Grass, Bermuda Grass, Johnson Grass, Para Grass, Russian Thistle, Canada Thistle, Bindweed, Whitetop (Hoary Cress) and many other grasses and weeds. Toxic to all types of vegetation. This material is recommended for vegetation control on non-cropped areas such as <u>ditch banks</u>, <u>fencerows</u>, along <u>roadsides</u>, around <u>buildings</u>, <u>power lines</u>, right-of-ways, and other similar waste areas.

ACTIVE INGREDIENTS:

Sodium Chlorate 6.00% Disodium Octaborate Anhydrous 11.00%

Caution: Keep Out of Reach of Children

See other precautions on the side panel.

EPA Reg. No. 11694-12

CONSTRUCTION CHEMICAL SPECIALTIES, INC. SHAWNEE MISSION, KANSAS 66203

CAUTION:

NON-SELECTIVE

..... 6.00% 11.00% 83.00%

of Children side panel.

KEEP OUT OF REACH OF CHILDREN

Avoid contamination of food and feed. Keep away from children and domestic animals. Do not contaminate water used for humans, domestic animal consumption or for irrigation.

Harmful if swallowed.

Do not apply to areas underlaid by roots of desirable trees and shrubs.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

NET CONTENTS: 55 gallons

