

FOR INDUSTRIAL AND INSTITUTIONAL USE ONLY

DIC-100 DISINFECTANT

Recommended for cleaning, disinfecting, and sanitizing in hospitals, insitutions, food plants, and commercial buildings.

ACTIVE INGREDIENTS:

Alpha-(p-Nonylphenyl)-omega-hydroxypoly (oxyethylene)-iodine complex (Providing 1.75% titratable iodine)	18.05%
Phosphoric Acid	16.00%

INERT INGREDIENT:	65.95%
Total	100.00%

Manufactured under the following U.S. Patents:	2,931,777
	2,977,315
	3,028,299


DIC-100 Disinfectant has its own built-in indicator of germicidal activity. When the amber color disappears, a fresh solution should be prepared. This product is not adversely affected by water hardness or low temperatures.

DANGER: KEEP OUT OF REACH OF CHILDREN


(See side panel for additional precautions and first aid)

EPA Registration No. 11694-38


NET CONTENTS: 1 GAL



Manufactured By
CONSTRUCTION CHEMICAL SPECIALTIES, INC.
 5747 Kessler
 Shawnee Mission, Kansas 66205



11694-38



DIRECTIONS FOR USE:

For general cleaning and disinfecting in one operation, use 3 ounces of DIC-100 DISINFECTANT to 5 gallons of water.

For porous or extremely dirty surfaces, use 6 ounces to 5 gallons of water

For sanitizing previously-cleaned surfaces, use 1 ounce of DIC-100 DISINFECTANT in 5 gallons of water.

HOSPITALS:

Use as recommended above in animal rooms and cages, cafeterias, morgues, laboratories, operating areas, patient rooms, bedframes, walls and floors, thermometers, instruments, rubber gloves, tubing & catheters. Allow at least 10 minutes contact time. Rinse all food-handling surfaces with potable water before reuse. Before disinfecting, instruments should be cleaned to remove all adhering blood and serous exudates.

Commercial Buildings, Institutions, and General Housekeeping:

Use 3 ounces of DIC-100 DISINFECTANT to 5 gallons of water for cleaning and disinfecting floors, walls, lavatories, shower rooms, refuse containers, kitchen areas, and lunch rooms. Rinse all food-handling equipment with potable water before reuse.

FOOD PLANTS:

To sanitize previously-cleaned food-processing equipment and utensils, use 1 ounce of DIC-100 DISINFECTANT to 5 gallons of water. Allow to drain but do not rinse. When used in this manner, DIC-100 DISINFECTANT does not require a potable water rinse, in accordance with the Federal Food Additive Regulation 121.2547.

DANGER:

Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician immediately.

NOTE TO PHYSICIAN:

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed