

[11694-48]

11694-48

DOUBLE ACTION CONTACT & RESIDUAL SPRAY

**FOR USE IN HOMES, MOTELS, INSTITUTIONS,
RESTAURANTS, AND FOOD PROCESSING
PLANTS.**

ACTIVE INGREDIENTS:

*+(5-Benzyl-3-furyl)methyl 2, 2-dimethyl-3-(2-methylpropenyl)cyclopropane-carboxylate	0.350% w/w
Related compounds	0.048%
Aromatic petroleum hydrocarbons	0.464%
INERT INGREDIENTS:	99.138%
	100.000% w/w

*Cis trans isomers ratio: max. 30% (±) cis and min. 70% (±) trans.

+U.S. Patent No. 3,465,007.

EPA Reg. No. 11694-48

EPA Est. 11694-KS-1

WARNING KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid contamination of feed and food-stuffs. Remove pets, birds and cover fish aquariums before spraying. Do not use on edible crops. Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

WARNING: Contents under pressure. Do not puncture. Do not expose container to temperature above 120°F. Never throw container into fire or incinerator.

NET WEIGHT: 18 OZ. (1 LB. 2 OZ.)

**DYMON, INC.
3401 KANSAS AVE.
KANSAS CITY, KANSAS 66106**

DOUBLE ACTION INSECT KILLER—This is a dual purpose formulation that can be used as a contact spray against flying and crawling insects and as a residual insecticide providing up to 3 or 4 weeks of control of roaches.

DIRECTIONS FOR USE

Shake well before each use. Remove protective cap. Aim spray opening away from person. Push button to spray.

HOUSEFLIES, MOSQUITOES, AND GNATS:

Indoors: Close all doors and windows. Point nozzle upward, direct the spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

TO REDUCE ANNOYANCE FROM FLYING INSECTS:

Outdoors: Reduce mosquitoes, gnats (midges), houseflies, biting flies, and flying moths (millers) by spraying their resting areas. Follow precautions listed above. Hold container 18 to 24 inches from shrubs, bushes, and grassy areas. Spray these areas with a slow sweeping motion at the rate of from 1 to 3 seconds per cubic feet of space while moving away from the treated areas. To reduce annoyance from these flying insects, treat open porches and patios in a similar manner. Spray only when the air is still. Allow spray to dissipate before occupying the treated area. Repeat as necessary.

ROACHES, EARWIGS, GRAIN MITES, SOWBUGS, SPIDERS, AND CENTIPEDES:

Contact Spray: Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, and storage areas. Spray directly on insects where possible. Repeat as necessary.

Residual Spray: Spray baseboards, cracks and crevices, around door and window frames, and other hiding places such as closets, in back of and under appliances, sinks and cupboards in the kitchen and bathroom. In basements, spray hiding places in cracks and crevices, lower walls, and dark places. Spray at the rate of 3 to 4 seconds per linear foot at a distance of 12 to 24 inches from the surface. Repeat as necessary.

WASPS: Applications should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places contacting as many insects as possible. Repeat as necessary.

ANTS: Apply to ant trails and around door sills, window frames and other places where ants may enter.

SHAKE WELL BEFORE USING