

October, 19-1975

11694-58

NOTICE

KEEP FROM FREEZ'NG. If this product is stored in cold or freezing weather it may separate or thicken. This will not harm the performance. Before using, warm to room temperature and stir thoroughly.

CAUTION

Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Flammable mixture — Keep away from heat or open flame.

NOTE

Injury may occur on certain ferns and some species of Crassula and Canalita Juniper. Do not reuse empty container. Destroy by burying. Also bury spillage and spray waste away from irrigation or domestic water supplies.

This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning equipment or disposal of wastes. Apply this product only as specified on this label.

KEEP OUT OF REACH OF CHILDREN

I.T.C. 20

ACTIVE INGREDIENTS:

Aromatic Petroleum Derivative
Solvents 30.00%
* * Malathion 55.00%

INERT INGREDIENTS: 15.00%

Nonylphenoxypoly (ethyleneoxy)
ethanol Calcium Doedecyl Benzene
Sulfonate

* * 0,0-Dimethyl Dithiophosphate of Diethyl Mercap-
tosuccinate

**CAUTION:
KEEP OUT OF REACH
OF CHILDREN**

(See side panel for other precautions)

DIRECTIONS

Japanese Beetle, White Grubs, White-fringed Beetle and Wire Worms on grass turf areas. Mix 1 gallon of Concentrate with sufficient water to cover one acre. After applying, wash into the soil. This will give control for one season. To give up to 10 years control, apply at the rate of 2 to 4 gallons of Concentrate in sufficient water to cover one acre.

Soil Web Worms on grass turf areas. Water thoroughly before applying. Apply Concentrate at the rate of 1 to 2 gallons in sufficient water to cover one acre. Do not water for several days after treating.

Aphids and Spider Mites on Ornamentals — Mix Concentrate at the rate of 1 1/2 pints per 100 gallons of water. Spray thoroughly with a wet coarse spray. Repeat when necessary.

When treating for aphids and spider mites, reapply every 2 or 3 weeks or when the insects reappear.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

EPA Reg. No. 11694-58
EPA Est. 11694-KS-1



KANSAS CITY, KANSAS 64106