

DIRECTIONS FOR USE (General Classification)

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product has been designed specifically for hospitals, food processing plants, dairies, restaurants, bars and other institutions where disinfection and sanitization is of prime importance.

DISINFECTION--To disinfect previously cleaned hard inanimate surfaces (such as walls, floors, table tops) apply product with mop, cloth, or sponge or mechanical sprayer so as to wet thoroughly. Allow to remain wet for 10 minutes and then let air dry.

DILUTION RATES General Disinfection--add 3 oz. per 5 gallons of water. Hospital Disinfection--add 3 1/2 oz. per 5 gallons of water.

For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into solution containing 1 1/2 oz. product per gallon of water for 10 minutes. For disinfection, a pre-cleaning step is required and recommended. Prepare a fresh solution for each use.

DEODORIZER--To deodorize, apply product as indicated under the heading "General Disinfection" above.

MILDEWSTAT--To control mold and mildew on pre-cleaned hard non-porous surfaces (such as floors, walls, table tops) add 3 oz. of product per gallon of water. Apply solution with a cloth, mop or sponge making sure to wet all surfaces completely. Let air dry. Prepare a fresh solution for each use. Repeat application at weekly intervals or when mildew growth reappears.

SANITIZATION--To sanitize previously cleaned and rinsed non-porous food contact surfaces, prepare 200ppm active quaternary solution by adding 1 oz. of product to 4 gallons of water.

To sanitize immobile items (such as tanks, chopping blocks, counter tops) flood the area with 200ppm product solution (1 oz. product diluted in 4 gallons of water) for at least 60 seconds making sure to wet all surfaces completely. Remove and let air dry. Prepare a fresh solution daily or more frequently as soil is apparent.

To sanitize mobile items (such as drinking glasses, eating utensils) immerse in 200ppm product solution (1 oz. of product diluted in 4

**CONTROL
SANITIZER
VIRUCIDE
DEODORIZER
DISINFECTANT**

JUL 06 1987
 U.S. Dept. of Agriculture
 Pesticide and Rodenticide Act
 as amended, for the pesticide
 registered under
 EPA Reg. No. 11694-84

ACTIVE INGREDIENTS	
n-Alkyl (60% C., 30% C., 5% C., 5% C.) dimethyl benzyl ammonium chlorides	5%
n-Alkyl (68% C., 32% C.) dimethyl ethylbenzyl ammonium chlorides	5
INERT INGREDIENTS	90
TOTAL	100

EPA REG. No. 11694-84
EPA EST. No. 11694-KS-1

KEEP OUT OF REACH OF CHILDREN

DANGER

See side panels for a full and complete primary and secondary treatment instructions.

NET CONTENTS



3401 KANSAS AVENUE KANSAS CITY, KANSAS 64106

DIRECTIONS (CONT)

gallons of water) for at least 60 seconds making sure to immerse completely. Remove and let air dry. Prepare a fresh solution daily or more frequently as soil is apparent.

This product provides sanitizing, disinfecting and deodorizing. Hospitals, Food Processing Plants, Dairies, Restaurants, Bars and other institutions. When used as recommended, it is effective as sanitizer against *Escherichia coli* and *Staphylococcus aureus*; and as disinfectant against *Staphylococcus aureus*, *Salmonella choleraesuis* and *Pseudomonas aeruginosa*. This product authorized under the Food Additive Amendment of the Federal Food, Drug and Cosmetic Act as a sanitizing solution at 200ppm (1 oz. of product diluted in 4 gallons of water) active level without requiring a final potable water rinse.

DISINFECTION--At 3 oz. per 5 gallon dilution, this product exhibits effective disinfectant activity against the organisms *Salmonella choleraesuis* and *Staphylococcus aureus*.

At 3 1/2 oz. per 5 gallons dilution, this product is an acceptable hospital disinfectant and demonstrates disinfectant activity against *Pseudomonas aeruginosa*.

VIRUCIDAL ACTIVITY--This product when used on any non-porous hard material surfaces at 3 1/2 oz. per 5 gallons of water is effective against Influenza A (representative of the flu virus), Herpes simplex (representative agent of blisters and mononucleosis), Adenovirus Type 1 (representative agent of upper respiratory infections) and Vaccinia (representative of the pox virus group).

Efficacy tests have demonstrated that this product is an effective bactericide and virucide in the presence of organic soils (5% brood serum). **DEODORIZATION**--This product deodorizes those areas which generally are hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, pet areas and any other areas which are prone to odors caused by microorganisms.

MOLD AND MILDEWSTAT--This product prevents and controls mold and mildew and the odors they cause on hard non-porous inanimate surfaces.

SANITIZATION--When used for sanitization of previously cleaned food equipment or food contact items (and active quaternaries at 200ppm) 1/2 product diluted in 4 gallons of water. At this level, a POTABLE WATER RINSE is required.

This product is an effective sanitizer when diluted in water at 200ppm (1/2 oz. per 5 gallons).

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, FEED OR STORAGE CONTAINERS.

DO NOT REUSE EMPTY CONTAINERS.

KEEP OUT OF REACH OF CHILDREN. For more information on safe use, see the label on the container.

CONTAINER DISPOSAL: Containers should be disposed of in accordance with local, state or federal regulations.

PRECAUTIONARY STATEMENTS
Hazard to Human and Domestic Animals
DANGER

Keep Out Of Reach Of Children. Causes severe eye and skin damage. Do not get in eyes, skin or on clothing. Harmful if swallowed. Wear goggles, face shield and rubber gloves when handling. Avoid contamination of food. Remove and wash contaminated clothing.

STATEMENT OF PRACTICAL TREATMENT
In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk and whole gelatin solution. If these are not available, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN
Probable mucosal damage may contraindicate the use of astringent lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

DIRECTIONS FOR USE (General Classification)

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product has been designed specifically for hospitals, food processing plants, dairies, restaurants, bars, and other institutions where disinfection and sanitization is of prime importance.

DISINFECTION-To disinfect previously cleaned hard inanimate surfaces (such as walls, floors, table tops) apply product with mop, cloth, or sponge or mechanical sprayer so as to wet thoroughly. Allow to remain wet for 10 minutes and then let air dry.

DILUTION RATES-General Disinfection--add 3 oz. per 5 gallons of water. Hospital Disinfection--add 3 1/2 oz. per 5 gallons of water.

For disinfection of previously cleaned surgical instruments, barber tools, and dental equipment, simply submerge instruments into solution containing 1 1/2 oz. product per gallon of water for 10 minutes. For disinfection, a pre-cleaning step is required and recommended. Prepare a fresh solution for each use.

DEODORIZE--To deodorize, apply product as indicated under the heading "General Disinfection" above.


MILDEWSTAT--To control mold and mildew on pre-cleaned, hard non-porous surfaces (such as floors, walls, table tops) add 3 oz. of product per gallon of water. Apply solution with a cloth, mop or sponge making sure to wet all surfaces completely. Let air dry. Prepare a fresh solution for each use. Repeat application at weekly intervals or when mildew growth reappears.

SANITIZATION--To sanitize previously cleaned and rinsed non-porous food contact surfaces, prepare 200ppm active quaternary solution by adding 1 oz. of product to 4 gallons of water.

To sanitize immobile items (such as tanks, chopping blocks, counter tops) flood the area with 200ppm product solution (1 oz. product diluted in 4 gallons of water) for at least 60 seconds making sure to wet all surfaces completely. Remove and let air dry. Prepare a fresh solution daily or more frequently as soil is apparent.

To sanitize mobile items (such as drinking glasses, eating utensils) immerse in 200ppm product solution (1 oz. of product diluted in 4

**CONTROL
SANITIZER
VIRUCIDE
DEODORIZER
DISINFECTANT**



ACTIVE INGREDIENTS	
n-Alkyl (60% C ₁₂ -30% C ₁₈ -5% C ₁₅ -C ₁₆) dimethyl benzyl ammonium chlorides	5%
n-Alkyl (68% C ₁₂ -32% C ₁₅ -C ₁₆) dimethyl ethylbenzyl ammonium chlorides	5%
INERT INGREDIENTS	90%
TOTAL	100%

EPA REG. No. 11694-84
EPA EST. No. 11694-KS-1

KEEP OUT OF REACH OF CHILDREN

DANGER

See side panel for additional precautions and practical treatment statements.

NET CONTENTS



3401 KANSAS AVENUE KANSAS CITY, KANSAS 64108

DIRECTIONS (CONT)

gallons of water for at least 60 seconds making sure to immerse completely. Remove and let air dry. Prepare a fresh solution daily or more frequently as soil is apparent.

This product provides sanitizing, disinfecting and deodorizing in Hospitals, Food Processing Plants, Dairies, Restaurants, Bars, and other institutions. When used as recommended, it is effective as a sanitizer against *Escherichia coli* and *Staphylococcus aureus*; and as a disinfectant against *Staphylococcus aureus*, *Salmonella choleraesuis* and *Pseudomonas aeruginosa*. This product authorized under the Food Additive Amendment of the Federal Food, Drug and Cosmetic Act for use as a sanitizing solution at 200ppm (1 1/2 oz. product diluted in 4 gallons of water) active level without requiring a final potable water rinse.

DISINFECTION-At 3 1/2 oz. per 5 gallon dilution, this product exhibits effective disinfectant activity against the organisms *Salmonella choleraesuis* and *Staphylococcus aureus*.

At 3 1/2 oz. per 5 gallons dilution, this product is an acceptable hospital disinfectant and demonstrates disinfectant activity against *Pseudomonas aeruginosa*.

VIRUCIDAL ACTIVITY-The product when used in recommended dilution on hard surfaces at 3 1/2 oz. per 5 gallons of water is effective against Influenza A (representative of the flu virus), Herpes simplex (causative agent of blisters and mouth sores), Adenovirus Type 4 (causative agent of upper respiratory infections), and Vaccinia Virus (representative of the pox virus group).

Efficacy tests have demonstrated that this product is an effective bactericide and virucide in the presence of organic soil (5% blood serum).

DEODORIZATION-This product deodorizes those areas which generally are hard to keep fresh smelling, such as garbage, storage areas, empty garbage bins and cans, pet areas and any other area which are prone to odors caused by microorganisms.

MOLD AND MILDEWSTAT-This product prevents and controls mold and mildew and the odors they cause on hard non-porous inanimate surfaces.

SANITIZATION-When used for sanitization of previously cleaned food equipment or food contact items in a active quaternary solution (1 oz. product diluted in 4 gallons of water). At this level, **POTABLE WATER RINSE** is required.

This product is an effective sanitizer when diluted in water at 1 1/2 oz. per 5 gallons CaCl₂.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

CONTAINER REUSE: Do not reuse.

Do not reuse empty containers.

REUSE OF CONTAINERS: Do not reuse containers for other purposes unless approved by the manufacturer.

CONTAINER DISPOSAL: Recycle where available. If not, burn in a secure incinerator.

PRECAUTIONARY STATEMENTS
Hazard to Human and Domestic Animals
DANGER

Keep Out Of Reach Of Children. Causes severe eye and skin irritation. Do not get in eyes, skin, or on clothing. Harmful if swallowed. Wear goggles, face shield and rubber gloves when handling. Avoid contamination of food. Remove and wash contact immediately.

STATEMENT OF PRACTICAL TREATMENT
In case of contact, immediately flush eyes with plenty of water for at least 15 minutes. If eyes are irritated, flush and wash contaminated clothing before reuse. If swallowed, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN
Probable mucousal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory distress, and convulsions may be needed.

STORAGE AND DISPOSAL
Do not contaminate water, food, or feed by storage or disposal.
CONTAINER REUSE: Do not reuse.
Do not reuse empty containers.
REUSE OF CONTAINERS: Do not reuse containers for other purposes unless approved by the manufacturer.
CONTAINER DISPOSAL: Recycle where available. If not, burn in a secure incinerator.