

11694-110

MASTER LABEL

All word(s), phrases, or symbols in { } and **bolded** are OPTIONAL

* Denotes Additional Brand Name

Page 1 of 2

THE END.

{*THE END. FLYING INSECT KILLER}

{*THE END. CRAWLING INSECT KILLER}

- {AUTHORIZED FOR USE IN NON-FOOD/FEEED AREAS OF FOOD/FEEED PROCESSING PLANTS AND OFFICIAL ESTABLISHMENTS OPERATING UNDER THE MEAT, POULTRY, SHELL EGG GRADING AND EGG PRODUCTS INSPECTION PROGRAMS.}
- {CONTAINS PYRENONE® AND PERMANONE®}
- {CONTROLS INDUSTRIAL INSECTS}
- {SEEKS OUT BUGS WHERE THEY HIDE}
- {KILLS ON CONTACT}
- {FOR COMMERCIAL, AGRICULTURAL/INSTITUTIONAL USE}
- {A BROAD SPECTRUM INSECTICIDE HIGHLY EFFECTIVE AGAINST {ANTS} {(INCLUDING CARPENTER ANTS),} {BEES,} {BEETLES} {(INCLUDING {CARPET,} {FLOUR,} {GRAIN,} {AND} {GROUND BEETLES}),} {CENTIPEDES,} {CLOVER MITES,} {COCK-ROACHES,} {CRICKETS} {(INCLUDING MOLE CRICKETS),} {WIGGS,} {FIREBRATS,} {FLEAS,} {MILLIPEDES,} {MOTHS,} {PILLBUGS,} {SCORPIONS,} {SILVERFISH,} {SOWBUGS,} {SPIDERS,} {TICKS,} {WASPS,} {AND OTHER PESTS LISTED ON THE LABEL}}
- {KILLS LICE AND THEIR EGGS}
- {FOR USE ON BEDDING, FURNITURE AND OTHER INANIMATE OBJECTS INFESTED WITH LICE}
- {NON-STAINING}
- {NO UNPLEASANT ODOR}
- {WATER-BASED FORMULA}
- {FOR USE ON PREMISES TO CONTROL FLEAS AND TICKS}
- {PROTECTS FOR UP TO {4 WEEKS,} {12 WEEKS,} {3 MONTHS}}

ACTIVE INGREDIENTS:

Pyrethrins	0.10%
*Piperonyl Butoxide, Technical	0.50%
**Permethrin	0.20%

OTHER INGREDIENTS: 99.20%
{100.00%}

*Equivalent to min. 0.4% (butylcarbityl)(6-propylpiperonyl) ether and 0.1% related compounds.

**{2-(4-chlorophenyl) methyl (+/-) cis/trans 3-(2,2-dichloroethenyl) 2, 2-dimethylcyclohexanecarboxylate. Cis/trans ratio: min. 35% (+/-) cis and max. 65% (+/-) trans. {Pyraperm, Pyrenone And Permanone - Registered Trademarks of Valent Biosciences Corp.}

EPA REG. NO. 11694-110 {-XXXX} EPA EST. NO. 11694-KS-1

{PRECAUCION AL CONSUMIDOR: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.}

KEEP OUT OF REACH OF CHILDREN

CAUTION


See {back, rear, side} panel for additional precautionary statements.

Net Weight XXXX



Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols

You can recycle this steel container in an increasing number of communities.



Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide Registered under EPA Reg. No. 11694-110

2/9/2004

1/2

PRECAUTIONARY STATEMENTS

Hazards To Humans & Domestic Animals

CAUTION. Harmful if inhaled. Avoid breathing vapor or spray mist. Remove contaminated clothing and wash before reuse.

FIRST AID

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to any body of water.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

STORAGE AND DISPOSAL

STORAGE: Store in a cool dry area away from heat or open flame. Do not contaminate water, food, or feed by storage and disposal.

DISPOSAL: Do Not puncture or incinerate!

If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use in food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where foods may be prepared or held.

In the home all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

Nonfood areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before spraying. Vacate room after treatment and ventilate before reoccupying. Do not allow children or pets to contact treated areas until surfaces are dry.

Do not use in rooms/areas while occupied by patients, the elderly or infirm. Do not apply to classrooms when in use.

SHAKE WELL BEFORE USING

Remove protective cap, hold container upright and spray from a distance of 12 to 15 inches.



{INDOORS:}

{CLUSTER FLIES,} {HOUSEFLIES,} {FRUIT FLIES,} {MOSQUITOES,} {GNATS,} {WASPS,} {HORNETS,} {BEES,} {BLACK FLIES,} {SKIPPER FLIES,} {BLOW FLIES} AND {SMALL FLYING MOTHS:} {Close all doors and windows and direct spray upward into center of room with a slow sweeping motion. Spray 5-10 seconds for average room, moving toward the door. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Leave room and keep area closed for 15 minutes after spraying. Ventilate room thoroughly before re-entry. Sweep up and discard fallen insects.}

{COCKROACHES,} {WATERBUGS,} {PALMETTO BUGS,} {SILVERFISH,} {FIREBRATS,} {CRICKETS,} {SCORPIONS,} {SPIDERS,} {MILLIPEDES,} {CENTIPEDES,} {SOWBUGS} AND {PILLBUGS:} {Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray when ever possible. }

{ANTS: Spray trails, nests and points of entry. Spray on ant where possible.}

{INDIAN MEAL MOTHS,} {ANGUMOIS GRAIN MOTHS:} {Close doors and windows, direct spray into all parts of room especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space, moving toward the door. Leave room and keep area closed for 15 minutes following treatment.}

{EXPOSED STAGES OF CONFUSED FLOUR BEETLES,} {CIGARETTE BEETLES,} {DRUG STORE BEETLES,} {SAW-TOOTHED GRAIN BEETLES,} {RED FLOUR BEETLES,} {RICE WEEVILS,} {GRANARY WEEVILS,} {GRAIN MITES IN EXPOSED ADULT AND LARVAE STAGES:} {Spray into cracks and crevices, processing and handling equipment, around cartons or containers or other areas where these insects tend to congregate. Contact as many insects as possible. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Space spray: Spray 20 seconds per 1000 cu. ft. of space, moving toward the door. Leave room and keep area closed for 15 minutes following treatment. Infested stored food products should be disposed of, or treated using an approved

{CARPET BEETLES: Spray directly on beetles, floors, baseboards, shelves and carpets.)

{BEDBUGS: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, molding and floors. Allow all sprayed articles to dry thoroughly before use.)

{TO CONTROL {CLOTHES MOTHS} AND {CARPET BEETLES} On wool and other proteinaceous fibers: Remove infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated by spraying lightly, keeping nozzle at least three feet from fabric to avoid staining.)

12/08/03
184-185MV2
KRH

PRN 2001-6
PRN 2001-1
PRN 1992-2

{FLEAS IN BUILDINGS - SPACE SPRAY: Hold can 36 inches above floor and direct spray toward floor and lower walls at a rate of 10 seconds per 100 sq. ft. making sure that all floor, sofa and chair surfaces are contacted. Leave room and keep area closed for 15 minutes. Open and ventilate before reoccupying. Repeat treatment after 7 days or as needed.)

{FLEAS & TICKS (ADULT AND LARVAE): Remove and thoroughly clean or destroy pet bedding. Spray pet sleeping quarters, bed, entire floor and floor covering, with special attention to cracks and crevices around baseboards, and window and door frames. Infested upholstered furniture and mattresses may also be treated with this product. A test application should be made to upholstery or drapery fabrics in an inconspicuous place before use.)

{TO KILL LICE AND LOUSE EGGS: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Hold container upright with nozzle away from you. Depress valve and spray from a distance of 8 to 10 inches. Spray each square foot for three seconds. Spray only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all sprayed articles to dry thoroughly before use.)

{ANIMAL QUARTERS AND MILK ROOMS: Remove animals prior to treatment. Rapid control of House Flies, Mosquitoes, Gnats, Wasps, Hornets, Bees and Small-Flying Moths, close windows and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.)

{TO TREAT LIVESTOCK: To protect beef cattle, dairy cattle, goats, sheep, hogs and horses from attacks of Stable Flies, Horse Flies, Deer Flies, Face Flies, House Flies, Horn Flies, Mosquitoes, and Gnats: Spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from Face Flies, spray the face and head, but do not spray into eyes. Repeat treatment when flies are troublesome.)

{FOR SWINE: Do not ship animals for slaughter within 5 days of last treatment.)

{TO CONTROL BLOOD SUCKING LICE: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.)

{TO CONTROL POULTRY LICE, spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.)

{FOR CONTROL OF BEDBUGS AND MITES IN POULTRY HOUSES, spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.)

{DOORS):

Hold spray nozzle away from face. Hold can on slight downward angle. For best results, spray when air is calm. Allow a few minutes for product to take effect. Spray with wind if breeze is blowing. Remove all exposed food and cooking utensils. Cover all food handling surfaces or wash thoroughly after treatment and before use. Do not use on edible crops.)

{FLYING INSECTS): {Flies,} {Mosquitoes,} {Gnats,} {Small Flying Moths,} {Hornets,} {Wasps} and {Yellow Jackets}. {Allow fog to drift over area. Direct spray toward bushes or grass from at least 4 to 6 feet away. Give special attention to breeding areas such as garbage cans and compost piles.)

{MUD DAUBERS AND WASP NESTS: Spray mist directly on insects and their nests from approximately 2 feet away. Contacted insects will fly away from fog. Applications should be made in late evening when insects are at rest. Spray into hiding and breeding places contacting as many insects as possible.)

{CRAWLING INSECTS): {Ants,} {Cockroaches,} {Including Asian Cockroaches,} {Crickets,} {Mole Crickets,} {Centipedes,} and {Silverfish}. {Hold can two (2) to three (3) feet from surface to be treated and spray infested surface of patio or picnic area, hitting as many insects as possible. Spray to slightly moisten surface. Also spray legs of tables and chairs to repel ants. Spray ant hills for best control.)

{Buyer assumes all risk of use, storage or handling of this product not in strict accordance with directions given herewith.)

{Manufactured, Distributed, Sold} (in the USA) (by, for)
ITW Dymon, Olathe, KS 66061

{For an MSDS, call XXX-XXX-XXXX or visit www.xxxxx.com and request document #xxxx}