

[11715-17]

SPEER

INDUSTRIAL INSECTICIDE

PROFESSIONAL STRENGTH

A PRESSURIZED, CONCENTRATED INSECTICIDE

FOR USE IN

FOOD PLANTS, RESTAURANTS, HOSPITALS,
FOOD MARKETS, HOTELS, MOTELS OR PLANTS

ACTIVE INGREDIENTS:

Pyrethrins	500%
*Piperonyl Butoxide, Technical	4.000%
Petroleum Distillates	8.000%

INERT INGREDIENTS: 87.500%

*Equivalent to 3.2% (Butylcarbityl) (6-Propylpiperonyl)
Ether and .8% of Related Compounds.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Contents under pressure. See back panel for other precautions.

Net Wt. 2.5 Lbs.

— DIRECTIONS —

FLIES, MOSQUITOS, GNATS, WASPS, FLYING ADULT MOTHS: Determine cubic feet of space to be treated. Example: 10' high x 10' long x 10' wide = 1000 cubic feet. Close all windows and doors. Direct valve opening away from person and spray from 2 to 5 seconds for each 1000 cubic feet of space. Disperse mist in all directions and, if possible, have fans operating to speed up circulation of insecticide throughout area being treated. Leave treated area. Keep area closed for at least 15 minutes after treatment. Ventilate room completely before reoccupying. Sweep up and destroy all insects.

RICE and GRANARY WEEVILS, CONFUSED FLOUR BEETLES, RED FLOUR BEETLES, CHOCOLATE MOTHS, DRUG STORE BEETLES in exposed adult and larvae stages: Contact as many insects as possible when treating infested areas such as cracks, crevices, machinery, and dark secluded areas. Spray 10 to 20 seconds per 1000 square feet of floor space. Repeat treatment weekly or as insects appear.

CARPET BEETLES: Make localized application to areas of the floor and baseboards. Spray directly into cracks and under carpets where insects may be found. Treat crevices, cracks or closets and infested areas of shelving. Repeat as necessary.

WARNING: KEEP OUT OF REACH OF CHILDREN

Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation.

In hospitals, patients should be removed from the room prior to treatment. Room should be ventilated for 2 hours after spraying. Do not return patients to room until after ventilation.

Avoid contamination of food and feedstuff. Cover or remove fishbowls. Remove all pets from area to be treated. In case of skin contact, wash with soap and water.

WARNING: Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130° F. may cause bursting. Never throw container into fire or incinerator.

TECHNICAL DATA — FLY CONTROL

Dosage required	1 sec. per 1000 cu. ft.
Treatment per 2.5 Lb. Container	1,135,000 cu. ft.
Delivery rate @ 70° F	1 gram per second
Cylinder	Aluminum
Coverage	Each 2.5 Lb. container will effectively treat approximately the same area as 17 gallons of an oil base spray without the use of over 16 gallons of petroleum distillate.

EPA Reg. No. 11715-17

Manufactured by
SPEER PRODUCTS, INC.
Memphis, Tennessee 38109