



# SPEER INDUSTRIAL INSECTICIDE

PROFESSIONAL STRENGTH

A PRESSURIZED, CONCENTRATED INSECTICIDE  
FOR USE IN

FOOD PLANTS, RESTAURANTS, HOSPITALS,  
FOOD MARKETS, HOTELS, MOTELS OR PLANTS

**ACTIVE INGREDIENTS:**

Pyrethrins	500%
*Piperonyl Butoxide, Technical	4.000%
Petroleum Distillates	8.000%

**INERT INGREDIENTS:** 87.500%

\*Equivalent to 3.2% (Butylcarbitol 16-Propyl piperonyl Ether and 8% of Related Compounds)

**KEEP OUT OF REACH OF CHILDREN.**

## CAUTION

Contents under pressure. See back panel for other precautions.

Net Wt. 2.5 Lbs.

Manufactured by:  
**SPEER PRODUCTS, INC.**  
MEMPHIS, TN 38118

EPA Reg. No. 11715-17 EPA Est. No. 11715-TN-1

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**FLIES, MOSQUITOS, GNATS, WASPS, FLYING ADULT MOTHS:** Determine cubic feet of space to be treated. Example: 10' high x 10' long x 10' wide = 1000 cubic feet. Close all windows and doors. Direct valve opening away from person and spray from 2 to 5 seconds for each 1000 cubic feet of space. Disperse mist in all directions and, if possible, have fans operating to speed up circulation of insecticide throughout area being treated. Leave treated area. Keep area closed for at least 15 minutes after treatment. Ventilate room completely before reoccupying. Sweep up and destroy all insects.

**RICE and GRANARY WEEVILS, CONFUSED FLOUR BEETLES, RED FLOUR BEETLES, CHOCOLATE MOTHS, DRUG STORE BEETLES** in exposed adult and larvae stages: Contact as many insects as possible when treating infested areas such as cracks, crevices, machinery, and dark secluded areas. Spray 10 to 20 seconds per 1000 square feet of floor space. Repeat treatment weekly or as insects appear.

**CARPET BEETLES:** Make localized application to areas of the floor and baseboards. Spray directly into cracks and under carpets where insects may be found. Treat crevices, cracks or closets and infested areas of shelving. Repeat as necessary.

## Precautionary Statements Hazards to Humans and Domestic Animals

### CAUTION

Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation.

In hospitals, patients should be removed from the room prior to treatment. Room should be ventilated for 2 hours after spraying. Do not return patients to room until after ventilation.

Avoid contamination of food and feedstuff. Cover or remove fishbowls. Remove all pets from area to be treated. In case of skin contact, wash with soap and water.

### PHYSICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130° F may cause bursting.

## STORAGE AND DISPOSAL

**Storage:** Store in a cool area away from heat or open flame.

**Disposal:** Do not reuse empty container. Wrap container and put in trash collection.