

SPEER

BIRD SPRAY

KILLS MITES & RED LICE

A special formulation for Canaries,
Parakeets and other caged birds.

ACTIVE INGREDIENTS:

Pyrethrins	0.09%
*Piperonyl Butoxide, Technical	0.18%
N-octyl bicycloheptene dicarboximide	0.30%
Petroleum Distillate	0.43%
Triethylene Glycol	0.10%
Propylene Glycol	0.10%

INERT INGREDIENTS: 98.80%
*Equivalent to .144% (butylcarbitol) (6-propylpiperonyl)
ether and .036% related compounds.

KEEP OUT OF REACH OF CHILDREN*
CAUTION
See rear panel for additional precautions.

MANUFACTURED BY
SPEER PRODUCTS, INC.
Memphis, Tennessee 38118

EPA Reg. No. 107-104

EPA Reg. No. 107-104

Net WL

BEST AVAILABLE COPY

STATEMENT OF PRACTICAL TREATMENT

IF INHALED- Remove victim to fresh air. Apply artificial respiration if indicated.
IF ON SKIN- Remove contaminated clothing. Wash skin with soap and warm water. Get medical attention if irritation persists.
IF IN EYES- Flush eyes with plenty of water. Obtain medical attention if irritation persists.
IF SWALLOWED- Call a physician or Poison Control Center immediately. Do not induce vomiting without medical supervision because of possible aspiration hazard.

Essential Safety Statements

Hazards to Human and Domestic Animals

CAUTION

Harmful if swallowed. Avoid breathing vapors. Wash hands with soap and warm water after use.

In the home, all food processing surfaces, exposed food and utensils... should be covered during treatment, or thoroughly washed before use.

Cover or remove fish aquariums before spraying.

Physical Hazards

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Storage: Store in a cool area away from heat or open flame.

Disposal: Do not re-use empty container. Wash container and put in trash collection.

"FOR THE OWNER WHO CARES"
SPEER BIRD SPRAY

Formulated especially to control Red Lice and Mites on Canaries, Parakeets and all Caged Birds.

DIRECTIONS

SHAKE WELL. Spray parakeets, canaries and other caged birds as follows:

FOR CAGE MITE CONTROL: Remove bird and thoroughly spray perches and cage. Clean and spray at regular intervals.

FOR BIRD LICE CONTROL: Leave bird in cage and spray directly on bird, holding container at least 3 feet from bird or any surface. Spray lightly - one burst of 2 or 3 seconds. Apply no more than 2 times per week.

ACCEPTED
JUL 2 1980
11715-20