PM- 10

′ ¶ ′

SPEER BIRD SPRAY

KILLS MITES & RED LICE

A special formulation for Canaries, Parakeets and other caged birds.

ACTIVE INGREDIENTS:

Pyrethrins	0.09%
Piperonyl Buloxide, Technical*	0.18%
N-octyl bicycloheptene dicarboximide	0.30%
Triethylene Glycol	0.10%
Propylene Glycol	0.10%
Propylene GlycolINERT INGREDIENTS**:	99.23%

*Equivalent to .144% (butylcarbityl) (6-propylpiperonyl) elher and .036% related compounds.

"Contains petroleum distillate

KEEP OUT OF REACH OF CHILDREN CAUTION

See rear panel for additional precautions

Manufactured by:
SPEER PRODUCTS, INC.
Memphis, Tennessee 38118

EPA Reg. No. 11715-20 EPA Est. No. 11715-TN-1

Net Wt.____

Contents under pressure. Do not use or store near heat of open flame. Do not puncture or incinerate container.

Exposure to temperatures above 130 F may cause bursting.

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidence for Phase 2 Response, Page 2.12, Active to thert Change in Status.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

STATEMENT OF PRACTICAL

TREATMENT

IF ON SKIN OR CLOTHING: Remove contaminated clothing. Wash skin with soap and warm water. Get medical attention if irritation persists.

IF IN EYES: Flush eyes with plenty of water. Obtain medical attention if irrilation persists.

IF SWALLOWED: Call a physician or Poison Control Center in mediately. Do not induce vomiting without medical supervision because of possible aspiration hazard.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Avoid breathing vapors. Wash hands with soap and warm water after use.

In the home, all food processing surfaces, exposed food and utensils should be covered during treatment, or thoroughly washed before use.

Cover or remove fish aquariums before spraying.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsis' int with its labeling.

STORAGE AND DISPOSAL

Storage: Store in a cool area away from heat or open flame.

Disposal: Do not re-use empty container. Wrap container and put in trash collection.

"FOR THE OWNER WHO CARES" SPEER BIRD SPRAY

Formulated especially to control Red Lice and Mites on Canaries, Parakeets and all Caged Birds.

DIRECTIONS

SHAKE WELL, Spray Parakeets, Canaries and other caged birds as follows:

FOR CAGE MITE CONTROL: Remove bird and thoroughly spray perches and cage. Clean and spray at regular intervals.

FOR BIRD LICE CONTROL: Leave bird in cage and spray directly on bird, holding container at least 3 feet from bird or any surface. Spray lightly - one burst of 2 or 3 seconds. Apply no more than 2 times per week.