

SPEER SCREW WORM SPRAY & WOUND PROTECTANT

CONTAINS KORLAN®


KILLS

Screw Worms, Blowfly Maggots and Ear Ticks

PROTECTS WOUNDS

From
Screw Worms and Blowfly Maggots

**FOR LIVESTOCK OF ALL AGES,
INCLUDING NEWBORN ANIMALS**



ACTIVE INGREDIENTS:

Ronnel* [0,0-dimethyl 0-(2, 4, 5-trichlorophenyl) phosphorothioate†]	2.5%
Xylene	0.5%

INERT INGREDIENTS97.0%

*Trademark of The Dow Chemical Company abroad.
†Prepared from technical grade.


**FOR EXTERNAL USE ONLY
LIVESTOCK REMEDY – NOT FOR HUMAN USE**

®Korlan is a Trademark of The Dow Chemical Company.

**CAUTION
KEEP OUT OF REACH OF CHILDREN**

Read carefully other precautions on back.

NET WT. 1 LB. (16 OZ.)



**LIVESTOCK REMEDY – NOT FOR HUMAN USE
FOR EXTERNAL USE ONLY**

(Blue Color added for marking treated areas)

— DIRECTIONS FOR USE —

FOR USE ON CATTLE, HORSES, SHEEP, GOATS AND SWINE: Hold container about 3 inches from wound or infested area. Be sure opening is directed toward area to be treated. Press valve down to spray. Wet thoroughly by spraying directly into and about 1 inch around wound.

SCREW WORM AND BLOWFLY INFESTATIONS: Use particular care to get spray into deep pockets caused by screw worms. Repeat applications at 5 to 7 day intervals until wound is healed.

TO PREVENT SCREW WORM AND BLOWFLY INFESTATIONS: Apply directly to all open or fresh wounds such as from dehorning, castration, shear cuts, etc. Treat navels of newborn animals. Freely bleeding wounds may require several treatments. Repeat applications at 5 to 7 day intervals until wound is healed.

EAR TICKS: Spray directly into animal's ear. Repeat applications at 5 to 7 day intervals as needed.

FLEECE WORM INFESTATIONS: Spray infested area liberally, working toward center of infestation. Shearing or removing the wool from the infested area is not necessary. Repeat application at 5 to 7 day intervals as needed.

**CAUTION
KEEP OUT OF REACH OF CHILDREN**

Harmful if swallowed. Injurious to eyes. Avoid contact with skin, eyes and clothing. In case of contact with human eyes, flush eyes immediately with plenty of water and get prompt medical attention. Avoid breathing spray mists. Wash hands thoroughly after using. Avoid contamination of feed and foodstuffs.

WARNING: Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Never throw container into fire or incinerator. Ronnel is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals. Atropine is an antidote. Consult a veterinarian at the first sign of an adverse reaction.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on label. Buyer assumes all risk of any such use and/or handling of this material when such use and/or handling is contrary to label instructions.

E.P.A. Reg. No. 11715-24 E.P.A. Est. 11715-TN-1

Manufactured by
SPEER PRODUCTS, INC.
Memphis, Tenn. 38109