



[11715-31]



Dairy Spray

FOR MILK HOUSES AND ANIMALS

**Kills House Flies, Horn Flies,
Stable Flies, Gnats and Roaches**

ACTIVE INGREDIENTS:

| | |
|--------------------------------|--------|
| Pyrethrins | .500% |
| *Piperonyl Butoxide, Technical | 4.000% |
| Petroleum Distillates | .800% |

INERT INGREDIENTS: 87.500%

*Equivalent to 3.2 (Butylcarbityl) (6 Propyl piperonyl) Ether and 8 of Related Compounds

CAUTION:

KEEP OUT OF REACH OF CHILDREN.

See left Panel for other precautions

Net Wt. 2.5 Lbs.

FOR HOME USE:

To kill *Roaches, Spiders and Silverfish* thoroughly spray insects and all possible hiding places such as cracks, baseboards, sinks, cabinets and shelves. Repeat treatment when necessary.

To kill *Wasps and Hornets* apply in late evening when all of the insects have returned to the nest. Spray liberally into hiding places and nest, contacting as many insects as possible.

CAUTION:

KEEP OUT OF REACH OF CHILDREN

CAUTION: Cover milking utensils and milk to prevent contamination from spray and dead or falling insects. Avoid contamination of feed and foodstuffs. When used in enclosed areas, do not remain in treated areas. Ventilate the area after treatment is completed.

Remove pets, birds, and cover fish aquariums before spraying.

WARNING:

Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130° F may cause bursting. Never throw container into fire or incinerator.

Manufactured by
SPEER PRODUCTS, INC.
Memphis, Tennessee 38109

EPA Reg. No. 11715-31
EPA Est. No. 11715 TN 1

A ready to use insecticide for control of stable flies, house flies, horn flies and gnats on lactating and nonlactating dairy cattle and beef cattle, when used as directed

DIRECTIONS

MILKING PARLOR AND MILK ROOM:

Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet

Stand in the center of the room. Direct fog upward and spray in all directions. Seconds may be counted by saying one thousand, two thousand, etc. Keep room closed for 15 minutes after treatment. Example: A room 20 feet long, 12 feet wide and 8 feet high requires 2 seconds if closed tightly. Otherwise spray for 4 seconds. Do not remain in treated areas. Ventilate the area after treatment is completed.

ANIMAL USE:

From approximately 24 inches, thoroughly spray entire animal's face as it is being released to pasture. Spot treat withers, shoulders and back where saliva accumulates from head tossing. Spray to wet the hair but not the skin.

STANCHION BARN USE:

Wear behind animal and direct DAIRY AEROSOL over the animal for approximately 1 second per animal. For most effective results, apply each morning.