



ACTIVE INGREDIENTS:

o-Isopropoxyphenyl
methylcarbamate*25%

INERT INGREDIENTS 99.75%

SENDRA

*BAYGON = Reg. T.M. of Parent Company of Farberbairn
Bayer GmbH, Leverkusen.

CAUTION

KEEP OUT OF REACH OF CHILDREN

See back panel for additional cautions.

NET WEIGHT 15 OZ.

SPEER

This ~~BAYGON~~ FLEA & TICK formulation provides long lasting killing power and protection for your pet.

DIRECTIONS

SHAKE WELL BEFORE USING. ALWAYS HOLD CAN UPRIGHT WHILE SPRAYING.

Spray the animal from a distance of 6 to 8 inches. Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and under the body. While spraying, fluff the hair so that the spray will penetrate to the skin.

Spray 20 to 30 seconds for a 25 pound animal. Spray Cats 5 seconds or less for a five pound animal. Repeat treatment as necessary but do not treat more than once every 7 days.

Spray the animal's living quarters and bedding thoroughly attempting to contact directly as many insects as possible. Spray into cracks and crevices where ticks may be found indoors. Allow about 15 minutes of ventilation before returning animal to area.

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing mist. Avoid contact with skin, eyes or clothing. Wash thoroughly after using. If illness should occur obtain prompt medical aid. Atropine is antidotal.

Do not spray on birds or around fishbowls. Do not spray into pet's eyes, face, scrotum or open wounds.

CAUTION

KEEP OUT OF REACH OF CHILDREN

WARNING: Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Never throw container into fire or incinerator.

Manufactured by:

SPEER PRODUCTS, INC.

Memphis, Tennessee 38109

EPA EST. NO. 11715-TN-1

EPA REG. NO. 11715-41