

SPEER FLEA & TICK SPRAY FOR DOGS & CATS

RAPID KILL OF FLEAS AND TICKS

ACTIVE INGREDIENTS:

o-Isopropoxyphenyl
methylcarbamate 0.25%

INERT INGREDIENTS 99.75%
..... 100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

See rear panel for additional precautions.

MANUFACTURED BY
SPEER PRODUCTS, INC.
Memphis, Tennessee 38118

EPA Reg. No. 11715-41

EPA Est. No. 1715-TN-1

Net Wt. _____

BEST AVAILABLE COPY

STATEMENT OF PRACTICAL TREATMENT

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.
IF ON SKIN: Remove contaminated clothing. Wash skin with soap and warm water. Get medical attention if irritation persists.
IF IN EYES: Flush eyes with plenty of water. Obtain medical attention if irritation persists.
IF SWALLOWED: Call a physician or Poison Control Center immediately. Do not induce vomiting without medical supervision because of possible aspiration hazard.

This PROPOXUR FLEA & TICK formulation provides long lasting killing power and protection for your pet.

PRECAUTIONARY STATEMENTS HAZARDS OF HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing mist. Avoid contact with skin, eyes or clothing. Wash thoroughly after using. If illness should occur obtain prompt medical aid. Atropine is antidotal. Do not spray on birds or around fishbowls. Do not spray in to pet's eyes, face, scrotum or open wounds.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Storage: Store in a cool dry area away from heat or open flame.
Disposal: Replace cap and discard container in trash. Do not incinerate or puncture.

Spray the animal from a distance of 6 to 8 inches. Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and under the body. While spraying, fluff the hair so that the spray will penetrate to the skin.

Spray 20 to 30 seconds for a 25 pound animal. Spray Cats 5 seconds or less for a five pound animal. Repeat treatment as necessary but do not treat more than once every 7 days.

Spray the animal's living quarters and bedding thoroughly attempting to contact directly as many insects as possible. Spray into cracks and crevices where ticks may be found indoors. Allow about 15 minutes of ventilation before returning animal to area.

Do not use on puppies and kittens less than four weeks old.

ACCEPTED
11 1980
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 11715-41