

Silverfish: Spray baseboards, window and door frames, book cases, storage areas, and other surfaces suspected of silverfish infestation. Repeat treatment as necessary.

Carpet Beetles: Spray directly on beetles, floors, baseboards, shelves, and carpets. Repeat treatment as necessary.

Bedbugs: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings, and floors. Repeat treatment as necessary.

Fleas: Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.

CAUTION: Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Vacate premises after treatment and ventilate before reoccupying.

Remove pets, birds and cover fish aquariums before spraying.

Apply this product only as specified on this label.

Avoid contamination of feed and foodstuffs. When using this product in food processing plants, foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when facility is not in operation.

WARNING: Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130°F. may cause bursting. Never throw container into fire or incinerator.

EPA Reg. No. 11715-44 EPA Est. No. 11715-TN-1

Manufactured By:
SPEER PRODUCTS, INC. Not Reviewed. Registrant claims to be in accordance with
 Memphis, Tennessee 38109 Reregistration of Pesticide Product Guidance for Phase 2
 Response, Page 2.12, Active to then Change in Status.

***SPEER**

INDUSTRIAL Pressurized Spray

FOOD PLANTS, RESTAURANTS, HOSPITALS, SCHOOLS, SUPERMARKETS, MOTELS, HOTELS, DAIRIES, and ANIMAL QUARTERS.

Provides Rapid Control of Flies, Mosquitoes, Gnats, Wasps, Small Flying Moths (Millers) and accessible stages of Roaches, Ants, Water Bugs, Silverfish, Flour Beetles and Weevils (named on back panel), Grain Mites, and Spiders.

ACTIVE INGREDIENTS:

Pyrethrins 0.5%
 *Piperonyl Butoxide, Technical 4.0%

**INERT INGREDIENTS: 95.5%

*Equivalent to 3.2% of (butylcarbityl) (6-propylpiperonyl) ether and 0.8% of related compounds.

**Contains petroleum distillates.

CAUTION:

KEEP OUT OF REACH OF CHILDREN

See side panel for additional precautions

NET WT. 2.5 LBS.

DIRECTIONS

SHAKE BEFORE USING AND AT INTERVALS DURING PROLONGED USE.

Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining of fabrics or surfaces, do not spray at close range.

Flies, Mosquitoes, Gnats, Wasps, and Small Flying Moths: Close all doors and windows and spray room at the rate of 1 second per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10 to 15 minutes after treatment. Repeat treatment as necessary.

Roaches, Water Bugs, Crickets, Spiders, Centipedes, and Sow Bugs: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains, and plumbing, hitting insects with spray when ever possible. Repeat treatment as necessary.

Ants: Spray trails, nests, and points of entry. Spray on ants where possible. Repeat as necessary.

Indian Meal Moth, Angoumois Grain Moth: Close doors and windows, direct spray into all parts of room especially around stored food product containers, pallets, and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed.

Confused Flour Beetles, Saw-Toothed Grain Beetles, Granary Weevils, Rice Weevils, Red Flour Beetles, Grain Mites: Spray into cracks and crevices, processing and handling equipment, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat applications as necessary.