

SPEER

INSECTICIDE DIAZINON®

(Contains 4 Lbs. Diazinon per gallon)

For Roaches, Ants, Brown Dog Ticks & Fleas

For Use Against Insects Resistant To Chlorinated Hydrocarbons

Meets Federal Specification O-I-520


Active Ingredients:

O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate* 47.18%

Aromatic petroleum derivative 30.03%

Inert Ingredients: 22.79%

*Diazinon® Trademark of Ciba-Geigy Corp. 100.00%



DIRECTIONS FOR USE.

ROACHES AND ANTS: Prepare a ½-percent emulsion by slowly adding 1¼ fluid ounces (2½ tablespoons) of concentrate to 1 gallon of water with constant stirring. Apply as a strip treatment to cracks, crevices and other roach harborages, and to ant runways. Use coarse, wet spray and apply until surface is wet to point of run-off (approximately 1 gallon per 1000 square feet). Reapply only when reinfestation is noted. Do not use for space spray or for residual treatments covering extensive indoor surfaces such as entire bulkheads and overheads. Preferably do not use in room occupied by children under 1 year of age.

FLY CONTROL (NOT INTENDED FOR LIVING QUARTERS):

"Spot" bait sprays. Apply a spray containing ½ pint of the concentrate and 1 pound of sugar in 2½ gallons of water with a knapsack or similar type sprayer to areas frequented by flies, such as doorways, around windows, and other areas where flies congregate. Repeat applications every 2 to 3 weeks as fly population warrants.

Liquid bait for sprinkling can application

Use 2 fluid ounces of the concentrate and 1 pound of sugar (2 cups of syrup or molasses) in 5 gallons of water and sprinkle lightly over the floor in areas frequented by flies. Burlap bags may be laid out in those areas frequented by flies and sprinkled with this mixture. Application every day or two will generally be required.

BROWN DOG TICKS AND FLEAS IN BUILDINGS

Prepare a ½ percent emulsion as for roaches and ants above. Thoroughly spray infested areas around baseboards, window and door frames, wall cracks, sleeping quarters of household pets, and localized areas of floors and floor coverings. Applications should be repeated as infestations warrant and as reinfestations occur. Fresh bedding should be placed in animal quarters following treatment. Do not treat animals with this formulation.

WARNING

KEEP OUT OF REACH OF CHILDREN

When used in dwellings, care should be taken to avoid deposits which could be frequently contacted by children. Care should also be taken to avoid spotting of wallpaper and fabrics. DO NOT CONTAMINATE FOOD, FOOD CONTAINERS, OR COOKING UTENSILS.

WARNING

Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

May be fatal if swallowed! May be absorbed through skin! Do not breathe spray mist. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contamination of feed or foodstuffs. Apply only under supervision of trained personnel. Use suitable protective clothing and devices when handling concentrate or when spraying extensively.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

For special applications, this may be diluted with diesel oil or kerosene. A slight haze may develop, but utility of insecticide will not be affected.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Net Contents: One U.S. Gallon
EPA Reg. No. 11715-51 EPA Est. No. 11715 TN 1
Manufactured By
SPEER PRODUCTS, INC.
Memphis, Tenn. 38109