

Reg # 11715-135 PM-10 1971

BACK YARD FARMER TOMATO & VEGETABLE INSECT SPRAY

Provides Rapid Control of Insects Listed on Side Panel
May be used on Food Crops to within One day of Harvest

ACTIVE INGREDIENTS:

Pyrethrins..... 0.200%

INERT INGREDIENTS *..... 99.800%

*Contains petroleum distillate

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional precautionary statements

Net Weight _____

Manufactured by:

SPEER PRODUCTS, INC.
Memphis, TN 38118

EPA Reg. No. 11715-135

EPA Est. No. 11715-TN-1

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of waste.

PHYSICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate con-

tainer. Exposures to temperatures above 130 F may cause bursting.

STATEMENT OF PRACTICAL TREATMENT

If inhaled: Remove victim to fresh air. Apply artificial respiration if indicated.

If on skin or clothing: Remove contaminated clothing. Wash skin with soap and warm water. Get medical attention if irritation persists.

If in eyes: Flush eyes with plenty of water. Obtain medical attention if irritation persists.

If swallowed: Call a physician or Poison Control Center immediately. Do not induce vomiting without medical supervision because of aspiration hazard.

STORAGE AND DISPOSAL

Storage: Store in a cool dry area away from heat or open flame.

Disposal: Replace cap and discard containers in trash. Do not incinerate or puncture.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read entire label and apply as directed.

SHAKE WELL BEFORE USING

For best results, use throughout the growing season to prevent buildup of insect populations. Spray plants from a distance of 3 1/2 to 4 feet using short bursts (one second). To prevent injury to plants, avoid visibly wetting the foliage.

Apply in the early morning or late evening when the air is still, treating all foliage with particular attention to the undersides of the leaves. Contact insects directly whenever possible. Repeat applications at 7 to 10 day intervals or after heavy rains. Do not apply to food crops within one day of harvest.

USE ON

Tomatoes

TO CONTROL

Flea Beetle, Hornworm, Colorado Potato Beetle, Stink Bug, Green Peach Aphid, Two Spotted Spider Mite, White Fly

Peppers,
Eggplants,

Colorado Potato Beetle, Leafhoppers

Potatoes

Blister Beetle, Green Peach Aphid, Flea Beetle

Cucumbers,
Squash

Striped and Spotted Cucumber Beetle,

Melons

Aphids, Two Spotted Spider Mite

Beans

Mexican Bean Beetle, White Fly, 12-Spotted Cucumber Beetle, Two Spotted Spider Mite

Sweet Corn

Japanese Beetle

Carrots


Leafhoppers

Peas

Aphids

Cabbage

Aphids, Cabbage Loopers



Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.