

SPI PYRENONE® FOOD PLANT INSECTICIDE

A Concentrated Pyrenone® Micro Emulsion

**CLEAR – STABLE – LEAVES NO OILY OR
OBJECTIONABLE RESIDUE**

Designed for Use in Edible Product Areas of Official Establishments Operating Under the Meat, Poultry, Shell Egg Grading and Egg Products Inspection Programs and in Other Food Processing Plants.

Apply with Conventional Mechanical or Compressed Air Equipment Designed and Adjusted to Deliver a True Aerosol Mist.

ACTIVE INGREDIENTS:

Pyrethrins 0.50%

Piperonyl Butoxide, Technical* 5.00%

INERT INGREDIENTS:** 94.50%
100.00%

*Equivalent to 4.0% (Butylcarbityl) (6 propyl piperonyl) ether and 1.0% related compounds

Pyrenone® - Registered trademark of Roussel Uclaf Corp.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panel for additional precautions.

NET CONTENTS _____

EPA Reg. No. 11715-154
EPA Est. No. 11715-TN-1

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION


Harmful if swallowed. Avoid contamination of feed and foodstuffs.

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not induce vomiting unless to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-757-4943 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply only as specified on this label.



11715-154

7/1/2003

1/2

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Close windows and doors, and shut off ventilating systems. Use a non-thermal fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

The above described application will kill accessible, exposed stages of:

FLIES, including FRUIT FLIES, MOSQUITOES, SILVERFISH, FLYING MOTHS, GNATS, CRICKETS, COCKROACHES, SPIDERS, SCORPIONS, WASPS, HORNETS, CLOVER MITES, CHEESE MITES, CHEESE SKIPPERS and these common stored product pests: GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES, SPIDER BEETLES, CIGARETTE BEETLES, DRUGSTORE BEETLES, ANGOUMOIS GRAIN MOTHS, BEDITERRANEAN FLOUR MOTHS, INDIAN MEAL MOTHS, TOBACCO MOTHS, MEAL MOTHS and GRAIN MITES.

In Mushroom Production and processing: To control adult MUSHROOM FLIES, FUNGUS GNATS and CECID FLIES in growing houses,

apply at a rate of 1 ounce per 1000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results refrigeration should be turned off. In packing and canning plants use at a rate of 1 ounce to 2000 cubic feet of space.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE AND SPILL PROCEDURES: Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and local procedures.

DO NOT TRANSPORT OR STORE UNDER 32°F.

[Distributors may choose their own emergency number and time of operation.]

**MANUFACTURED BY:
SPEER PRODUCTS
4242 B. F. GOODRICH BLVD.
MEMPHIS, TN 38118**

Buyer assumes all risk of use, storage or handling of this product not in strict accordance with directions given herewith.

Netserver/Reg./SPILabels11715-154-NOTIF2001-1&6(3-20-03)