KILLS & REPELS - Fleas, Ticks, Lice, Stable Flies, Hom Flies, Houseflies, Face Flies, Horse Flies, Deer Flies, Bat Flies, Mosquitoes and Gnats.

#### PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Avoid inhalation of vapors. Avoid contact with skin, eyes or clothing. If in eyes, flush with plenty of water. Call a physician If imitation persists. Wash skin with warm water and soap. Do not contaminate food or foodstuffs.

Avoid treatment of nursing puppies or kittens. If treatment is necessary, spray fingertips and rub into animal's coat. Do not soray into eyes, face, or genitals. If reaction should occur, contact your veterinarian immediately.

This product is toxic to fish. Keep out of takes, streams, ponds, tidal marshes and estuaries.

#### PHYSICAL HAZARDS

Flammable. Keep away from heat and open flame. Do not smoke white spraying.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Do not use or store near heat or open flame. Store in a dry area inaccessible to children.

Disposal: Do not reuse empty bottle. Rinse thoroughly before discarding <u>in Irash.</u>

ACCE PANILE EDIHOUGH LICENSED VETERINARIANS

MANUFACTURED BY OCT 1 6 1987 SPEER PRODUCTS, INC. MEMPHIS, TN 38118

Under ju l'adoral Insactivide, Pungicido, oud Redontioide Act, an amended, for the pesticide fegiatored under

🗰 A 🌬 . No.

# SPEER

(Liquid)

# FLEA & TICK **KILLER**

With Repellent and Deodorant

KILLS Fleas, Ticks and Lice

REPELS

Gnats, House Flies, Stable Flies and Mosquitoes

**DEODORIZES & CONDITIONS** 

### **ACTIVE INGREDIENTS:**

Pytethrins	0.150% w/
*Piperonyl butoxide, technical	
N-Octyl bicycloneptene	
Gicarboximide	0.500%
2,3,4,5-Bis(2-butylene) tetrahydro-	
2-furaldehyde	0.500%
INERT INGREDIENTS	97.350%
<del></del>	80.00

\*Equivalent to 1.20% (butylcarbityl) (6 propyl piperonyl) Bither and 0.30% other related compounds.

FOR USE ON DOGS, CATS, PUPPIES, KITTENS, HORSES, AND PONIES.

# KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional precautions.

Net Contents \_\_\_\_\_

EPA Res No 11715-160

EPA ENT NO 11715-TN-1

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

## ĬŬŜE ONLY IN WELL VENTILATED AREA

CAUTION: The product may squirt out of the nozzie when the protective nozzle cap is removed—if the temperature is higher than room temperature. Remove protective nozzle cap from sprayer by rotating tab to up position and pressing away from nozzle. Retain protective cap and re-attach between uses. Spray at a distance of 6-12 inches by squeezing trigger repeatedly and rapidly.

### DOGS AND CATS

Cover animal's eves with hand and with a firm fast stroke spray head, ears and chest until damp. With fingertips rub into face and around mouth, nose and eyes. Then spray neck, middle and hind quarters, finish legs last.

For best results, direct spray against the natural lay of the hair. On long haired dogs, rub your hand against the lay of the hair, spraying the ruffled hair directly behind the hand. Make sure spray thoroughly wets ticks. Repeat treatment as needed.

### **PUPPIES AND KITTENS**

Spray back of pet only, then spray on your lingertips and rub into animal's coat and around mouth. nose and eyes.

#### KENNELS OR PET SLEEPING QUARTERS

Apply thoroughly to infested areas such as pet beds and resting quarters, nearby cracks and crevices, along and behind baseboards, window and door sitis, and localized areas of floor and floor covering where fleas, ticks, and tice may be present. Old pet bedding should be removed and replaced with clean fresh bedding after treatment of pet areas. Treat pets with this product before allowing them to enter treated area. Do not spray surfaces where spotting, staining, or discoloration would be objectionable

### HORSES AND PONIES

To control stable flies, horse flies, deer flies and face flies, remove excess did from legs, flanks, top line and other body areas commonly attacked by these flies. To treat face, dampen a cloth with spray and rub on face and head, being careful to avoid eyes. Repeal treatment as necessary.